

Climate Change and Health

CO2 COALITION





Climate Change and Health



CO2 COALITION

CO2Coalition.org



Table of Contents

Introduction	2
Warmth, Wealth and Health	2
Figure 1: The Preston curve of Life Expectancy at birth increasing with GDP per capita	3
Figure 2: Projected economic impact of global warming of 1°C, 2.5°C and 3°C over the 21st century	4
Temperature and Disease	4
Figure 3: Nocturnal temperature trends for 1910-39 (Top) and 1970-97 (Bottom): trends plotted for the coldest nights from the left (No. 1) to the warmest (No. 365)	5
Figure 4: Distribution of cholera in 2016 as reported to the World Health Organisation.....	7
Figure 5: Projected changes in cold-related and heat-related excess mortality over the 21st century for nine regions and three emissions scenarios: RCP2.6, RCP4.5 and RCP8.5	9
Disease Vectors	10
Figure 6: Changing global malaria endemicity 1900-2007	11
Figure 7: Global distribution of Aedes aegypti in 2015	13
Figure 8: Distribution of Aedes albopictus as of December 2007 CDC	13
Figure 9: Progress of WNV in the U.S. 1999-2003. White 0, Blue <1%, Green 1-5%, Yellow 5-10%, Red >10%	15
Figure 10: WNV cases reported to the CDC 2014-2016. White 0, Green light 1-10, dark 21-50, Black >50	16
Figure 11: West Nile virus neuro-invasive disease incidence reported to CDC by year, 1999-2016	16
Figure 12: Reported cases of Lyme disease in the United States in 2016	18
Figure 13: Reported cases of Lyme disease in the U.S., 1996-2016	18
Figure 14: Vector and reservoir data points in North America (2009 data).....	20
Extreme Weather Events	21
Figure 15: Annual Number of Major Atlantic Basin Hurricanes, 1730-2005: Reconstructed from coral and plankton sediments in the Caribbean Sea and Observed.	21
Figure 16: Strength Index of Tropical Cyclone Events in North Qld., 1226-2003	22
Figure 17: Number of severe and non-severe tropical cyclones in Australia, 1970-2017.....	22
Figure 18: Annual number of strong tornadoes across the United States, 1954-2014.....	24
Figure 19: Brisbane River heights in metres, 1840-2016	25
Figure 20: Australian flood fatality rate from 1900 to 2015.....	25
Figure 21: Annual mean rainfall across Australia, 1900-2017.....	26
Figure 22: Flood fatalities in the continental United States, 1959-2005.....	26
Figure 23: Colorado River stream flow, Observed: 1905-2005 & Reconstructed from Tree Rings: 762-2005.....	27
Figure 24: Global tropospheric temperature anomalies, June-August 2003	28
Figure 25: Mean Temperature Deciles for the 2016-17 Australian Summer.....	29
Figure 26: Rank of the June–September 1954 heat wave based on National Climatic Data Center (NCDC) climate division temperature data for the years 1895–2009	30
Food, Famine, Climate and CO2	33
Figure 27: Global population (billions) and mortality from great famines (millions).....	34
Figure 28: Global wheat production this century (thousand tons pa)	35
Social Strife and Mental Health	37
Energy Sources and Health	38
Figure 29: Prevalence of pneumoconiosis among U.S. underground coal miners and NSW coal workers.....	40
Table 1: Pounds of CO2 emitted per million British thermal units (Btu) of energy for various fuels	41
Figure 30: Material resources (tonnes) per unit energy produced (TWh)	43
Table 2: Mortality rate per unit of power produced by energy sources (2012).....	45
Conclusion	45
References.....	46



Introduction

Human health, morbidity, mortality and longevity are significantly impacted by climate. This review examines the evidence for past, present and possible future health impacts of climate change on disease and disease vectors, on extreme weather events, floods, droughts, heatwaves and wildfires, on food and famine, and on social and mental health. It will also examine health issues in relation to various energy sources and conclude with an overview of priorities.

Warmth, Wealth and Health

For over two million years, Earth has been in the grip of an Ice Age interspersed every 100,000 years or so with interglacial warm periods lasting about 10,000 years. While the present warming began about 18,000 years ago, the Holocene interglacial dates from 11,700 years ago when South Greenland warmed 7°C in just 50 years.¹ The human population exploded in a warmer world. Agriculture began and civilisations arose during the Holocene Climatic Optimum,² globally warmer than now³ for several millennia in Russia⁴ and up to 7°C warmer than now in the Arctic.⁵ Subsequent gradual cooling has been interrupted by the Minoan, Roman and Medieval warm periods, during which mankind multiplied and flourished.

When rapid cooling after the Roman Warm Period ushered in the Dark Ages, the bubonic Plague of Justinian (541-542CE) killed 13% of the world's population, 25 million people, and twice that over the next two cold centuries.⁶ Civilisation again flourished during the Medieval Warm Period,⁷ after which a cold and miserable Little Ice Age (LIA) resulted in frequent widespread crop failures, mass starvation, disease and depopulation.⁸ Crop failures over successive summers from 1315 produced the Great Famine of Europe⁹ with "extreme levels of crime, disease, mass death and even cannibalism."¹⁰ The Black Death (1346-1353) wiped out 30-60% of Europe's population and up to 200 million people across Eurasia.¹¹

Global rewarming since the 18th century, plus better housing, sanitation, food and water supplies has greatly benefited human health and prosperity. Deaths from typhoid and tuberculosis declined dramatically during pre-antibiotic 20th century warming (1910-1945).¹² As temperatures rose, mortality from all causes fell.¹³ From a billion people in 1800, the global population had doubled by 1927, and again to four billion in 1974. It reached six billion in 2000 and is now 7.6 billion people.¹⁴

Northern winters have nevertheless remained more lethal than summers.¹⁵ In the United States, winters were 13% more lethal in 1952-1967 and even more so (16%) despite global warming in 1985-1990, largely due to the increased availability and affordability of air-conditioning.¹⁶ Despite warming from 1964 to 1998, Davis *et al.* (2003)¹⁷ found a 74.4% decline in heat-related mortality in 28 of the largest U.S. cities. They estimated that another 1°C increase would further reduce the net mortality rate.¹⁸ Analysing over 74 million deaths in 384 locations across 13 countries, Gasparrini *et al.* (2015)¹⁹ found that cold weather was twenty times more lethal than hot weather. About 7.7% (3% to 11%) of all deaths were related to non-optimal temperatures, nearly 7.3% to cold and only 0.42% to heat. The 2017-18 UK winter death-toll was the worst in 42 years,²⁰ due largely to fuel poverty²¹ affecting almost a third of the elderly.²²



Life expectancy is strongly associated with wealth or GDP per capita (Fig. 1).²³ Both have been increasing in all regions of a warming world since the industrial revolution (1770-2015).²⁴




Figure 1: The Preston curve of Life Expectancy at birth increasing with GDP per capita

Source: https://en.wikipedia.org/wiki/Preston_curve

Fossil fuels have cheaply and reliably powered industry, mechanised agriculture and transport, helped to end slavery and emancipate women and children, propelled urbanisation with its sewerage and safe water supplies, provided electricity for heating, refrigeration and air-conditioning, and facilitated better hospitals and health care. Fossil fuel products and by-products (fertilisers, pesticides and carbon dioxide) also boosted food production and human nutrition. Fossil fuels also contributed to global warming; by how much no one can say precisely, but probably about half earlier estimates.²⁵ While it is possible to have too much of a good thing, we are a long way from doing that globally. Richard Tol (2010)²⁶ analysed fourteen peer-reviewed papers examining the likely impacts on GDP and human welfare of warming this century – by 1°C (2 studies), 2.5°C (10 studies) and 3°C (2 studies) – and found that that another 1-2°C would probably increase global GDP (Fig. 2). As shown in our White Paper, **A Primer on Carbon Dioxide and Climate**,²⁷ global warming this century is unlikely to exceed 1°C, even without climate action. Dire predictions are based on flawed models, unhealthy exaggerations and a failure to factor in human ingenuity.²⁸




Figure 2: Projected economic impact of global warming of 1°C, 2.5°C and 3°C over the 21st century.
Source: Tol, 2010, derived from UN data

Temperature and Disease

By absorbing some solar radiation by day and emitting it to the surface by night, greenhouse gases moderate the **diurnal temperature range (DTR)**.²⁹ Whereas solar warming increases daytime maximum temperatures, greenhouse gases primarily impact nocturnal minimums. Knappenberger *et al.* (2001)³⁰ nicely demonstrated this in a study of night-time temperatures across the U.S. during 1910-1939 and 1970-1997. Whereas the first warming period, due primarily to solar **influences**, showed a fairly even distribution across the year, the latter warming period impacted primarily the coldest nights (Fig. 3). Cold nights have been decreasing and warm nights increasing right across the globe.³¹ Temperature variability has thus reduced.³²

High temperature variability is associated with increased cardiovascular and respiratory mortality in the United States.³³ For every 1°C increase in DTR, cardiac mortality increases by 1.7% in Hong Kong³⁴ and by up to 3.2% in Shanghai.³⁵ Emergency visits to the Huashan Hospital (Shanghai) with respiratory infections were 1% and 2% higher for every 1°C increase in the current-day and 2-day moving average DTR respectively.³⁶ Increasing atmospheric CO₂ actually **reduces** DTR.




Figure 3: Nocturnal temperature trends for 1910-39 (Top) and 1970-97 (Bottom): trends plotted for the coldest nights from the left (No. 1) to the warmest (No. 365).

Source: Knappenberger, Michaels and Davis, 2001

Asthma, now affecting over 300 million children,³⁷ has been increasing by 50% per decade in many countries.³⁸ The Centers for Disease Control (CDC) linked³⁹ this to climate change, but alternative explanations include increasing hygiene,⁴⁰ antibiotic use⁴¹ and the pasteurisation⁴² of cow's milk.⁴³ Rising CO₂ concentrations and temperatures may increase ragweed pollen numbers and perhaps other pollens associated with respiratory allergies,⁴⁴ but ragweed pollen allergenicity can vary four-fold,⁴⁵ and pollen numbers are spatially and temporally highly variable.⁴⁶ Ragweed pollen has actually been decreasing in



Zurich since 1982, and the major allergenic pollens have been declining in Basel, Switzerland since the early 1990s, as has the incidence of hay fever.⁴⁷ The Poaceae family of herbaceous grasses affects 80% of pollen allergy sufferers in Europe.⁴⁸ Jato *et al.* (2009)⁴⁹ found that the Poaceae pollen count had declined in four Spanish cities since 1993, by about 75% in Lugo and 80% in Santiago. They also found a delayed onset and shorter duration of the atmospheric pollen season. Xu *et al.* (2013)⁵⁰ found that childhood asthma increased 0-9 days after a DTR above 10°C, and also a 31% increase in emergency department admissions per 5°C increment in DTR. By reducing the DTR, CO₂ may actually be reducing the incidence of asthma.

Eczema and associated itch is more common in winter due to drying of the skin. A study of atopic eczema in the mountainous area of Davos, Switzerland in 1983–1989 found itch-intensity to be inversely correlated with temperature.⁵¹ Thirty Norwegian children improved in severity of eczema, quality of life, skin bacterial culture and medication usage after spending a month in the Canary Islands, and the improvement compared to a matched control group was still apparent three months later.⁵²

Gastroenteritis, which caused about 1.3 million deaths globally in 2015,⁵³ is predicted to increase with global warming.⁵⁴ Xu *et al.* (2013)⁵⁵ found that even in Brisbane, Queensland, “a 1°C increase in diurnal temperature range was associated with a 3% increase of Emergency Department Admissions for childhood diarrhea.” By reducing DTR, CO₂ may actually be reducing the global incidence of severe gastroenteritis.

Cholera is now confined to developing countries in the tropics and subtropics (Fig. 4) where it afflicts 3-5 million people and kills about 100,000 annually.⁵⁶ After spreading from Europe to London in 1831, an epidemic wiped out 50,000 people in just over a year. When another epidemic broke out in 1848, Dr John Snow⁵⁷ performed the world’s first epidemiological studies, linking it to contaminated water. Nearly a century and a half later, a paper in the prestigious journal *Science*⁵⁸ linked a 1991 outbreak in South America to climate change. The real cause, however, was a failure of the Peruvian authorities to properly chlorinate water supplies.⁵⁹ Climate change can be a handy scapegoat for government failure!

Respiratory disease is strongly related to temperature. Coughs, colds and influenza are far more prevalent in winter. Consultations for respiratory disease in London (1992-1995) increased linearly by 10.5% per degree (below 5°C).⁶⁰ Keatinge and Donaldson (2001)⁶¹ found a linear increase in mortality from respiratory disease in Londoners aged over 50 as the temperature fell below 15°C. The bronchiolitis season in central England was shortened by about three weeks for every 1°C of annual warming from 1981 to 2004.⁶² Carder *et al.* (2005)⁶³ analysed non-violent deaths in Scotland (1981-2001) and found that every 1°C drop in the daytime mean temperature below 11°C on any one day was associated with a 4.8% increased respiratory mortality over the following month. Respiratory-related deaths were




Figure 4: Distribution of cholera in 2016 as reported to the World Health Organisation.
Source: [http://gamapserver.who.int/mapLibrary/Files/Maps/Global_Cholera\(WER\)_2016.png](http://gamapserver.who.int/mapLibrary/Files/Maps/Global_Cholera(WER)_2016.png)

47% higher in winter than in summer in Oslo during the period 1990-1995.⁶⁴ Even in Sao Paulo, Brazil, Gouveia *et al.* (2003)⁶⁵ found the fewest respiratory deaths in all age groups occurred at 20°C; mortality increased twice as much per degree below 20°C as it did above 20°C. DTR is a risk factor for chronic obstructive pulmonary disease (COPD) mortality.⁶⁶

Cardiovascular disease (CVD) is the major cause of death worldwide. It presents more often in winter and is more often fatal on cold days.⁶⁷ A study in the Hunter Region of New South Wales, Australia (1985-1990) found that “fatal coronary events and non-fatal definite myocardial infarction were 20–40% more common in winter and spring than at other times of year”; coronary deaths were up to 40% more likely to occur on cold days than at moderate temperatures.⁶⁸ Cardiovascular mortality was 15% higher in the colder months October-March than in April-September in Norway (1990-1995),⁶⁹ 33% higher in Californian winters (1985-1996)⁷⁰ and 50% higher in mid-winter than in mid-summer in both London (1994-1996)⁷¹ and Israel (1976-1985),⁷² despite summer temperatures often exceeding 30°C.





Braga *et al.* (2002)⁷³ compared cardiovascular mortality in “hot” cities in the southern United States with “cold” cities in northern states. They found neither hot nor cold weather had much impact in the “hot” cities but significantly increased the mortality in the “cold” cities, where the cold-day effect was five times as great as the hot-day effect and persisted for days. There was a deficit of deaths for a few days after the hot days, indicating that hot days had a “harvesting effect” on those who were about to die. Cagle and Hubbard (2005)⁷⁴ examined the relationship between temperature and out-of-hospital cardiac deaths in people over 54 years of age in King County, Washington (USA) over the period 1980-2000; mortality rose by 15% on days with maximum temperatures below 5°C and dropped by 3% on days with maximum temperatures over 30°C. In Japan (1970-1999), heart disease and stroke mortality rates were nearly twice as high in winter as in summer.⁷⁵

Stroke is more common in cold weather. Novosibirsk, Siberia has one of the world’s highest rates of stroke, 87% being ischaemic (due to blocked cerebral blood vessels) and 32% higher on days with low ambient temperature.⁷⁶ A similar association was found in Korea, where Hong *et al.* (2003)⁷⁷ found a 24-48-hour lag between exposure to cold and the onset of stroke. A study of World Health Organisation (WHO) data on women aged 15-49 from 17 countries in Africa, Asia, Europe, Latin America, and the Caribbean found that a 5°C reduction in mean air temperature was associated with a 7% increase in hospitalisation with stroke.⁷⁸ Aneurismal subarachnoid haemorrhage is also strongly correlated with winter and cold weather.⁷⁹

Future Mortality: Keatinge and Donaldson (2004)⁸⁰ report: “even in climates as warm as southern Europe or North Carolina [USA], cold weather causes more deaths than hot weather. . . global warming will reduce this at first [but] the improvement is not likely to continue without action to promote defences against cold.” They conclude: “the overall effect of global warming on health can be expected to be a beneficial one.” Evaluating the future impact of unchecked global warming on human health, Bosello, Roson and Tol (2006)⁸¹ projected 1.4 million fewer deaths annually to 2050 and a lower mortality rate until at least 2200.

The Lancet, however, published a paper by Gasparrini *et al.* (2017)⁸² indicating that without mitigation, heat-related mortality will overtake cold-related mortality in Southeast Asia very soon, in Central and South America by mid-century, and in Southern Europe and overall before 2100. Using the worst emissions scenario (RCP8.5), they projected temperature increases ranging from 3.3°C for Australia to 4.9°C for North America, and 4.1°C globally, above the 2010-19 mean. The RCP4.5 scenario, projecting a mean global increase of 1.7°C, resulted in cold remaining more lethal than heat except in SE Asia and South America (Fig. 5). Russia, Alaska, Africa, the Middle East, northern and central Asia, India and Indonesia were ignored in this study, as was **adaptation and demographic changes**. The authors thus caution: “The reported figures should therefore be interpreted as potential impacts under well defined but hypothetical scenarios, and not as predictions of future excess mortality.”




Figure 5: Projected changes in cold-related and heat-related excess mortality over the 21st century for nine regions and three emissions scenarios: RCP2.6, RCP4.5 and RCP8.5

People are not potted plants – they take steps to reduce risk. Oren Cass⁸³ cautions: “If a projection of climate-change cost ignores **adaptation**, we can safely ignore it.” When Christidis *et al.* (2010)⁸⁴ factored adaptation into an analysis of the benefit of global warming on the mortality rate in England and Wales during the period 1976-2005, they found the lives-saved to lives-lost ratio increased from 29.4 to a phenomenal 121.4.

Recent **demographic changes** favour warmth: from 2006 to 2016, Australia’s hottest capital city, Darwin (latitude 12°S) had the nation’s fastest population growth (29%).⁸⁵ Southerners flock to sunny Queensland every winter, and many stay. So many U.S. citizens move from the cold northeast states to warm southwest states that it contributed 3-7% of the late 20th century gains in longevity, and is thought to delay about 4,600 deaths annually.⁸⁶ People are still voting with their feet on climate change. It is religious, political and tribal conflict that has been driving migration in Africa, the Middle East and Myanmar. We see no signs of Singaporeans, Malaysians or Indonesians fleeing the heat.



Disease Vectors

Vector-borne diseases⁸⁷ account for over 17% of all infectious diseases, causing more than 700,000 deaths annually. Vector-borne infection is regarded by the IPCC as a major climate change challenge to human health.⁸⁸ It is postulated that global warming will indirectly impact human health by spreading the following disease vectors to areas that have been too cool for them.⁸⁹

Vector	Species	Diseases
Mosquitoes	<i>Anopheles</i> <i>Aedes</i> <i>Culex</i>	Malaria – <i>P. falciparum</i> , <i>P. vivax</i> , <i>P. ovale</i> and <i>P. malariae</i> <i>Chikangunya</i> , <i>Dengue</i> , <i>Yellow Fever</i> , <i>Zika</i> West Nile Virus
Ticks	<i>Ixodes</i>	Tick-borne encephalitis, Lyme disease
Sandflies		Leishmaniasis

Malaria is caused by protozoan parasites belonging to the Plasmodium type, transmitted by the female Anopheles mosquito. It was endemic for millennia in Europe and England;⁹⁰ in areas near England’s marshes, 17th century burial records reveal a mortality rate comparable to that in sub-Saharan Africa today.⁹¹ Apart from epidemics during the unusually hot summers of 1848 and 1859, there was a near-linear decline in endemic malaria in the UK as the climate warmed from 1840 to 1910. It disappeared in 1953, and re-establishment with further warming is considered highly unlikely.⁹² Endemic malaria in Finland likewise faded out over two centuries of warming with limited or no counter measures or medication, leading Hulden and Hulden (2009)⁹³ to conclude that, “malaria in Finland basically was a sociological disease and that malaria trends were strongly linked to changes in the human household size and housing standard.” Helsinki had its last malaria epidemic in 1902, but parts of northern Europe including the Arctic Circle had devastating epidemics until the middle of the 20th century.⁹⁴

A malaria epidemic affected 30% of the population of the Tennessee River Valley in 1933.⁹⁵ During a period of rapid warming from 1916 to 1937, there was a four-fold decline in deaths from malaria in Mississippi, the only significant correlation being with family income: the higher the income the fewer the deaths.⁹⁶ Malaria remained endemic in 36 states until the CDC was created to tackle it after World War II. Swamps were drained, agricultural practices were changed, cases were isolated from mosquitoes and treated, mosquito nets and DDT were widely used and millions of homes were sprayed. The U.S. was considered malaria-free in 1951.⁹⁷ During another period of rapid warming (1977-2001), the incidence of malaria in northern Thailand declined by 6.45% per year, from 41.5 to just 6.72 cases per 100,000 people.⁹⁸ The global mortality rate per capita has declined by 95.4% since 1900.⁹⁹ The 20th century also saw a dramatic decline in malarial endemicity globally (Fig. 6).

Africa now bears the highest infection burden of any continent, with nearly 200 million cases reported in 2006. Malaria kills 0.7 to 2.7 million people every year, 90% of them in Africa, and three out four are children. Studies extending over 10-32 years found no correlation between temperature or rainfall and malaria incidence in western Africa,¹⁰⁰ or at four highland sites in east Africa¹⁰¹ or in western Kenya.¹⁰² A comprehensive study across Africa over an 85-year period (1911-1995) found a correlation



with precipitation in Southern Mozambique but none anywhere with temperature.¹⁰³ In a review paper, Rogers and Randolph (2006)¹⁰⁴ attribute the observed increase in malaria in many parts of Africa to “land-cover and land-use changes and, most importantly, drug resistance rather than any effect of climate,” noting “the recrudescence of malaria in the tea estates near Kericho, Kenya, in East Africa, where temperature has not changed significantly, shows all the signs of a disease that has escaped drug control following the evolution of chloroquine resistance by the malarial parasite.”




Figure 6: Changing global malaria endemicity 1900-2007. Source: Gething *et al.* 2010¹⁰⁵

- a, Pre-intervention endemicity (approximately 1900)
- b, Contemporary endemicity for 2007 based on a recent global project to define the limits and intensity of current *P.falciparum* transmission.
- c, Change in endemicity class between 1900 and 2007. Negative values denote a reduction in endemicity.



Haque *et al.* (2010)¹⁰⁶ analysed monthly malaria case data for the malaria endemic district of Chittagong Hill Tracts in Bangladesh from January 1989 to December 2008. They found no correlation with temperature, rainfall or humidity, but a strong negative association with the normalized difference vegetation index (NDVI), a satellite-derived measure of surface vegetation greenness. They state: “each 0.1 increase in monthly NDVI was associated with a 30.4% decrease in malaria cases” probably due to increasing insectivorous bird populations. By greening the Earth,¹⁰⁷ CO₂ may actually be reducing malaria risk.

It is thought that global warming will allow malaria to climb to higher altitudes.¹⁰⁸ Before the introduction of DDT and other public health measures, however, malaria transmission occurred much higher than now, at up to 2,600m in Kenya, 2,450m in Ethiopia, 2,500m in the Himalayas, 2,180m in Argentina and 2,773m (near thermal springs) in Bolivia.¹⁰⁹ After considering forest clearance, agriculture, urbanisation, health infrastructure, drug and insecticide resistance, civil strife and other influences on endemic malaria in the Highlands of Kenya and New Guinea, Paul Reiter of the Insects and Infectious Disease Unit of the Institut Pasteur in Paris, France concluded: “simplistic reasoning on the future prevalence of malaria is ill-founded; malaria is not limited by climate in most temperate regions, nor in the tropics, and in nearly all cases, ‘new’ malaria at high altitudes is well below the maximum altitudinal limits for transmission.” He also states “obsessive emphasis on ‘global warming’ as a dominant parameter is indefensible; the principal determinants are linked to ecological and societal change, politics and economics.”

When GDP per capita was included in models of future global malaria distribution, climate change proved to be much weaker.¹¹⁰ Simplistic models project an increase with climate change, but one using five variables and a high-emission scenario actually produced a 0.92% **decrease**. Economist, Indur Goklany (2004)¹¹¹ calculated that the malaria death toll could be halved through a combination of proven measures for a tiny fraction of the cost of mitigation.

Chikungunya virus (CHIKV) was first isolated on the Makonde Plateau of Tanzania during an outbreak in 1953.¹¹² The name is Makonde for “contorted or bent up” with incapacitating arthralgia. The primary host in Africa are non-human primates and small mammals, and its primary vector is the mosquito *Aedes aegypti*. CHIKV has been spreading rapidly in recent decades but not due to climate change. *A. aegypti* escaped Africa centuries ago with the slave trade and established itself widely across tropical and subtropical regions (Fig. 7), even in southern Europe until the mid-1900s when it disappeared during a period of warming.¹¹³ It recently re-established itself at Madeira, Portugal¹¹⁴ and in the Caucasian region bordering the Black Sea.¹¹⁵




Figure 7: Global distribution¹¹⁶ of *Aedes aegypti* in 2015. Source: Moritz UG Kraemer *et al*¹¹⁷

CHIKV broke out on Reunion Island and neighbouring Indian Ocean islands in 2005 and in India in 2006. It then spread to South-East Asia,¹¹⁸ simultaneously undergoing a genomic micro-evolution which enabled it to be transmitted by *Aedes albopictus*, the tiger mosquito of SE Asia.¹¹⁹ *A. albopictus*, which is intolerant to extreme heat,¹²⁰ has been adapting to cold climates in temperate regions such as Japan. It can hibernate over winter and produce eggs that are more cold-tolerant.¹²¹ Adult mosquitoes can even survive freezing winters in suitable microhabitats.¹²² It spread to Albania in 1979¹²³ and to Genoa, Italy in imported used tyres in 1990.¹²⁴ By 2007, it had spread extensively across southern Europe (Fig. 8). Ravenna in northern Italy experienced Europe's first CHIKV epidemic after the virus was introduced from India.¹²⁵ *A. albopictus* has recently spread westward in the continental United States, where locally acquired cases occurred in 2014,¹²⁶ but only travel-associated cases have been reported since 2015.¹²⁷




Figure 8: Distribution of *Aedes albopictus* as of December 2007 CDC¹²⁸



Modelling by Fischer *et al.* (2013)¹²⁹ projected an increased risk “for Western Europe (e.g. France and Benelux-States) in the first half of the 21st century and from mid-century onwards for central parts of Europe (e.g. Germany). Interestingly, the southernmost parts of Europe do not generally provide suitable conditions in these projections.” In other words, their emissions scenarios will make it too warm for *A. albopictus* to remain established in southern Europe. The reality, of course, is that no one knows what might happen this century; warming may be less than thought, vaccines and public health measures may control or even eradicate the virus from the developed world, it may mutate again, or more CO₂ may even reduce the incidence. Tuchman *et al.* (2003)¹³⁰ grew the quaking aspen (*Populus tremuloides*) (Michaux) trees at atmospheric CO₂ concentrations of either 360ppm or 720ppm for an entire growing season and fed the incubated leaf litter to four species of mosquito larvae to assess the effect on development. They found the larvae of *Aedes albopictus* had a mortality rate 2.2 times higher when fed the high-CO₂ litter, which delayed the development of all larvae by 9-20 days. When there are so many variables, who can predict the future!

Dengue fever is arguably the most important vector-borne viral disease globally, infecting over 200 million people, 1% of them severely with over 20 thousand deaths annually.^{131, 132, 133} A review of 16 studies, carefully selected from 75 papers with methodologies of varying quality, found dengue transmission to be highly sensitive to climatic conditions, especially temperature, rainfall and relative humidity.¹³⁴ Dire predictions¹³⁵ are usually based on the following facts: there has been a 30-fold increase over the past 5-6 decades; dengue is now endemic in 119 countries; the transmission zone has expanded to include half the world's population; further warming will lengthen the lifespan of the mosquito and shorten the extrinsic incubation period of the dengue virus, resulting in more infected mosquitoes for a longer period of time.¹³⁶ This must be weighed against other historical facts, scientific evidence and opinion:

- Endemic dengue has actually diminished or disappeared from some countries. In the United States, dengue and yellow fever were major public health problems during the cold 17th century but disappeared during the warm 20th century.¹³⁷ The CDC states: “As recently as the 1940s, large dengue outbreaks were documented in the United States reaching places as far north as Boston. Today, the situation has changed significantly. Reasonable climate, competent mosquito vectors, and susceptible human hosts are all still present in the continental United States, and dengue viruses are frequently reintroduced by infected travelers.¹³⁸ Transmission in the U.S. is rare, however, because there is insufficient contact between infected humans, vector mosquito species, and susceptible humans to sustain transmission.” *Aedes aegypti* and dengue were once prevalent in many parts of Australia, including New South Wales. As Russell *et al.* (2009)¹³⁹ point out, the current absence “is not because of a lack of a favourable climate”, nor is a temperature rise of a few degrees “likely to be responsible for substantial increases in the southern distribution of *A. aegypti* or dengue, as has been recently proposed.” Russell states: “of itself, climate change as currently projected is not likely to provide great cause for public health concern with mosquito-borne disease in Australia.”¹⁴⁰



- There are alternative explanations for the observed global increase in incidence: rapid urbanisation and international travel. Wilder-Smith and Gubler (2008)¹⁴¹ also note that “the disruption of vector control programs, be it for reasons of political and social unrest or scientific reservations about the safety of DDT, has contributed to the resurgence of dengue around the world.” They conclude: “population dynamics and viral evolution offer the most parsimonious explanation for the observed epidemic cycles of the disease, far more than climatic factors.”
- In 1995, dengue afflicted over 4,000 Mexicans in Tamaulipas while Texas, just across the border, had only a handful of non-imported cases. The essential difference was not climate but living standards and sound public health policies.¹⁴² Laredo, Texas (population 200,000) and Nuevo Laredo, Tamaulipas (population 290,000) are connected by bridges across the Rio Grande. After an outbreak in 1999, Reiter *et al.* (2003)¹⁴³ found “the incidence of recent cases, indicated by immunoglobulin M antibody serosurvey, was [12.3 times] higher in Nuevo Laredo, although the vector, *Aedes aegypti*, was more abundant in Laredo.” Reiter *et al.* determined that “the proportion of dengue infections attributable to lack of air-conditioning in Nuevo Laredo was 55% . . . [and] if the current warming trend in world climates continues, air-conditioning may become even more prevalent in the United States, in which case, the probability of dengue transmission will likely decrease.” The CDC¹⁴⁴ endorsed this: “Studies on the U.S.-Mexico border, for example, suggest that the restriction of transmission there is due to the limitation of contact between human hosts and mosquito vectors that comes with low housing density and the use of air conditioning and screens.”
- Future solutions to dengue could include the development of effective dengue vaccines,¹⁴⁵ genetic modification of *A. aegypti*,¹⁴⁶ the sterile insect technique¹⁴⁷ (shown to reduce the target mosquito population by more than 90%) and the *Wolbachia* bacterium¹⁴⁸ which allows mosquitoes to be resistant to arboviruses such as dengue and Zika.

West Nile Virus (WNV) was first identified in a West Nile district of Uganda in 1937. It is asymptomatic in 80% of infected people but can cause severe encephalitis or meningitis in about 1 in 150 infected persons, especially the elderly or immuneocompromised. It is transmitted by a *Culex* species of mosquito that has bitten an infected bird (not human). Appearing in New York in 1999 and spreading across the states taking hundreds of lives, it was soon linked to climate change.^{149 150} But its rapid spread from northeast to the south and west (Fig. 9) and its decline despite warming (Figs. 10 and 11) indicates that the vector was already there and climate change had nothing to do with that.




Figure 9: Progress of WNV in the U.S. 1999-2003. White 0, Blue <1%, Green 1-5%, Yellow 5-10%, Red >10%




Figure 10: WNV cases reported to the CDC 2014-2016. White 0, Green light 1-10, dark 21-50, Black >50




Figure 11: West Nile virus neuro-invasive disease incidence reported to CDC by year, 1999-2016

Source of Figs 9-11: CDC Cumulative Maps and Data¹⁵¹ and Wikipedia¹⁵²

Yellow fever spread from Africa to the Americas and Caribbean via the slave trade. The first recorded outbreak was in 1647 on the island of Barbados.¹⁵³ It reached New York City in 1668 and Philadelphia a year later; a 1793 epidemic in Philadelphia,¹⁵⁴ then the national capital, wiped out nearly 10% of its population.¹⁵⁵ Major outbreaks hit New Orleans in 1833 and 1853, and Memphis in 1878.¹⁵⁶ It spread to Europe early in the 19th century and took the lives of thousands in Gibraltar¹⁵⁷ and Barcelona.¹⁵⁸ There were even small outbreaks in France¹⁵⁹ and Wales.¹⁶⁰ Thanks to an effective vaccine¹⁶¹ and stringent travel regulations, it is now confined to central Africa and South America. The story of yellow fever illustrates that science and public health policies far outweigh risks associated with global warming.



Zika virus was first identified in the 1960s in South-East Asia, where it produced nothing more than a mild illness: fever, rash and aching joints. In 2013, a Zika strain suddenly appeared in French Polynesia, then in the Caribbean and Brazil, where it exploded and resulted in over 4,000 cases of microcephaly between late 2015 and early 2016. By September 2017, there were thought to have been between three and four million cases of Zika across 84 countries, due not to “unprecedented warming” but to international air travel. A Singaporean outbreak that began in August 2016 resulted in 455 cases over three months. When Singapore’s Ministry of Health and National Environment Agency quickly identified and managed infected people, eradicated mosquitoes and removed breeding sites, new cases were reduced by 48 percent within a month.¹⁶² Public health measures can effectively control such outbreaks before climate mitigation gets its boots on! If climate action impedes proven preventive measures, it will be counterproductive.

Tick-borne encephalitis virus (TBEV), another flavivirus, is the most significant vector-borne disease in Eurasia, with a mortality rate of 1-2% and up to 24% in the Far East. Randolph and Rogers (2000)¹⁶³ studied the effect of warming on the rodent-tick vector and concluded: “fears for increased extent of risk from TBEV caused by global climate change appear to be unfounded.” Because rapid cooling in autumn allows more larvae and nymphs to feed together in the following spring, the short TBE viraemia making this important for pathogen transmission, Randolph (2001)¹⁶⁴ suggested that TBEV transmission should be negatively affected by a warmer climate. Reported cases in Estonia and Lithuania¹⁶⁵ declined during the warm 1998 El Niño and were even lower in 1999. The end of Soviet rule and subsequent socioeconomic transition produced conditions favourable for transmission.¹⁶⁶ Randolph concluded: “there is increasing evidence from detailed analyses that rapid changes in the incidence of tick-borne diseases are driven as much, if not more, by human behavior that determines exposure to infected ticks than by tick population biology that determines the abundance of infected ticks . . . the evidence is that climate change has not been the most significant factor driving the recent temporal patterns in the epidemiology of tick-borne diseases.”¹⁶⁷

Lyme disease was first diagnosed in 1975 in Old Lyme, Connecticut,¹⁶⁸ from whence it derived its name. It is the most common tick-borne human disease, with an estimated annual incidence of 300,000 in the United States¹⁶⁹ and at least 85,000 in Europe.¹⁷⁰ It is caused by the spirochete bacteria, *Borrelia burgdorferi* and sometimes by *Borrelia mayonii*.¹⁷¹ It is transmitted in the eastern United States and parts of Canada by the tick, *Ixodes scapularis*, and on the Pacific Coast by *I. pacificus*.¹⁷² As these ticks like habitats with at least 85% humidity and need temperatures over 7 °C (45 °F) during host questing in spring,¹⁷³ the northeast United States is especially suitable (Fig. 12).




Figure 12: Reported cases of Lyme disease in the United States in 2016. Source: CDC Lyme Resources Brochure¹⁷⁴

The impact of climate change on Lyme disease appears to be complex. Subak (2003)¹⁷⁵ found a correlation between warmer winters and the incidence of Lyme disease the next summer, perhaps because mild winters enhance survival of the ticks' primary host, the white-footed mouse. Warm dry summers, on the other hand, are associated with a reduced incidence, perhaps because of reduced survival of both mouse and *Ixodes* nymph.¹⁷⁶ The number of reported cases of Lyme disease appears to have peaked in 2009, despite further warming (Fig. 13).




Figure 13: Reported cases of Lyme disease in the U.S., 1996-2016. Source: CDC Lyme Resources Brochure¹⁷⁷



Modelling by Brownstein *et al.* (2005):¹⁷⁸ “generated the current pattern of *I. scapularis* across North America with an accuracy of 89% ($P < 0.0001$). Extrapolation of the model revealed a significant expansion of *I. scapularis* north into Canada with an increase in suitable habitat of 213% by the 2080s. Climate change will also result in a retraction of the vector from the southern U.S. and movement into the central U.S.” As their modelled transmission zone migrates northwards into Canada over the next 70 years, it retreats from Florida and Texas, and the population exposed to Lyme actually diminishes, by 28% in the 2020s, by 12.7% in the 2050s and by 1.9% in the 2080s. The connection between suitable *I. scapularis* and deciduous forest is so strong that the authors state: “recent emergence of Lyme disease throughout the northeastern and mid-Atlantic states has been linked to reforestation”. The motor car may thus have contributed to the emergence of Lyme disease by converting numerous redundant horse-paddocks into woodlands and by fertilising them with carbon dioxide.

In Europe and Asia, the vectors of Lyme borreliosis (LB) are *I. ricinus* (Europe) and *I. persulcatus*.¹⁷⁹ Like its North American cousin, *I. ricinus* prefers forest to open land and deciduous to conifer.¹⁸⁰ Late 20th century warming has been linked to ticks spreading into higher latitudes and altitudes (observed in the Czech Republic¹⁸¹) and to higher incidences of LB,¹⁸² though distorted by better reporting over time in most regions. Moreover, the incidence of LB actually declined after 1995 in the Czech Republic and Lithuania.¹⁸³ In a 2006 WHO publication¹⁸⁴ on Lyme borreliosis in Europe, Elisabet Lindgren and Thomas G.T. Jaenson state:

The impact of climate variability on disease occurrence is more complex than the effects that the climate may have on the distribution and population density of the vectors. Studies on the effects of long-term climate variations on tick-borne disease prevalence are affected by several scientific problems. Data vary between places and over time owing to differences in awareness, surveillance, and diagnostic methods, which so far have made longterm studies on LB difficult to perform.

As with TBEV, the impact of climate change on the vectors and transmission of Lyme disease is complex and uncertain. The focus should be on educating the public, early diagnosis and treatment rather than climate mitigation.

Leishmaniasis, from over 20 *Leishmania* species of protozoa parasite, is transmitted by an infected female sandfly, with over 50 species of the genus *Phlebotomus* in the Old World and genus *Lutzomyia* in the New World. The main animal reservoirs include rodents, dogs, wild cats, jackals, foxes, sloths, hyraxes, and other carnivores. It is endemic in 88 countries across Africa, Asia, Europe, and North and South America, infecting up to a million people and killing up to 30,000 annually, mainly among the poorest people on Earth.¹⁸⁵ It is associated with malnutrition, population displacement, poor housing, a weak immune system and lack of financial resources. It is also linked to environmental changes such as deforestation, building of dams, irrigation schemes, and urbanization.



The disease manifests in three forms: visceral, cutaneous, and mucocutaneous.¹⁸⁶ The most common form is cutaneous leishmaniasis which occurs in the Americas, the Mediterranean basin, central Asia and the Middle East, and has been increasing in Egypt.¹⁸⁷ Visceral leishmaniasis (kala-azar) is highly endemic in Sudan, is strongly associated with malnutrition and fatal in 95% of untreated cases. It is thought to have spread from there to the Indian subcontinent and the New World.¹⁸⁸ Over 90% of mucocutaneous leishmaniasis occurs in Ethiopia, Bolivia, Brazil and Peru. In North America, leishmaniasis is endemic in Mexico and Texas and has begun to expand its range northward (Fig. 14). Models indicate that climate change may extend the risk of human exposure northwards, even into parts of southern Canada.¹⁸⁹




Figure 14: Vector and reservoir data points in North America (2009 data).
A: Both vector species. B: All four reservoir species. Source: Gonzalez *et al.* (2010)¹⁹⁰

Such modelling, of course, is as uncertain as the climate modelling on which it is based. It also assumes that the United States will be no more effective than African nations in preventing the spread of this disease and its vectors.

Extreme Weather Events

The deadliest natural disasters include hurricanes/typhoons/cyclones, storms/tornadoes, floods, droughts, heatwaves and wildfires. It is claimed that anthropogenic climate change is making these worse and thus putting more lives at risk.

Cyclones in the Southern Hemisphere are the same as North Atlantic hurricanes and Northwest Pacific typhoons except that they spin clockwise. Wind (120-300+ km/h), storm surge and flooding rain make these one of the deadliest natural disasters. A tropical cyclone in 1970¹⁹¹ killed an estimated 300,000 people in what is now Bangladesh. The deadliest Atlantic hurricane was probably the Great Hurricane of 1780,¹⁹² which took about 22,000 lives. Major hurricanes hit New York City in 1815¹⁹³ and 1821,¹⁹⁴ and another in 1893.¹⁹⁵ Atlantic hurricane activity actually declined from 1800 and was unusually quiet during the period 1970-1995 (Fig. 15). In 2013, no major hurricane made landfall for the first time in 45 years, and hurricanes were the fewest in number since 1982.




Figure 15: Annual Number of Major Atlantic Basin Hurricanes, 1730-2005: Reconstructed from coral and plankton sediments in the Caribbean Sea and Observed. Source: Nyberg *et al.* (2005)¹⁹⁶

Water vapour reaching high altitudes and precipitating as cyclonic rain contains a higher proportion of oxygen-16 isotope, which is lighter than oxygen-18. Nott *et al.* (2007)¹⁹⁷ used this to analyse cyclonic activity over 800 years from stalagmites in a Chillagoe cave in North Queensland and found that the most intense cyclones preceded white settlement (Fig. 16). The frequency and intensity of Australian cyclones since 1970 is shown in Figure 17.




Figure 16: Strength Index of Tropical Cyclone Events in North Qld., 1226-2003
Source: Nott *et al.* 2007.




Figure 17: Number of severe and non-severe tropical cyclones in Australia, 1970-2017. Severe tropical cyclones are those with a minimum central pressure less than 970 hPa. Source: Bureau of Meteorology¹⁹⁸

Haig and Nott (2016)¹⁹⁹ later developed “a new tropical cyclone activity index spanning the last 1500 years” and found “that in addition to other well-known climate indices, solar forcing largely drives decadal, interdecadal, and centennial cycles within the tropical cyclone record.”

Cyclones have caused over 2,100 deaths in Australia since 1839.²⁰⁰ Category 5 Cyclone Mahina²⁰¹ was Australia’s deadliest natural disaster, killing over 300 people in 1899. Thanks to better forecast and warning systems, and to improved building codes in cyclone-prone regions, loss of life from cyclonic activity is now rare – not a single death in 2011 from Category 5 Cyclone Yasi.²⁰² While deaths have



dramatically declined, damages have increased, but not due to climate change. According to the World Meteorological Organization,²⁰³ “the recent increase in societal impact from tropical cyclones has largely been caused by rising concentrations of population and infrastructure in coastal regions.” After normalizing the mainland U.S. hurricane damage during 1900-2005 to 2005-values, Pielke *et al.* (2008)²⁰⁴ found no trend in damages. They were notably low during the 1970s and 1980s. The 1926 Miami hurricane²⁰⁵ caused the most damage: \$157 billion normalized.

Typhoons²⁰⁶ account for nearly a third of the world’s tropical cyclones. The deadliest was Typhoon Nina which took nearly 100,000 lives in China when flooding caused 12 reservoirs to fail in 1975.²⁰⁷ The most intense were Typhoon Tip in 1979,²⁰⁸ with wind speeds of up to 310km/h before making landfall in the central Philippines, and Typhoon Haiyan in 2013,²⁰⁹ with wind speeds up to 314km/h. The most active typhoon season was in 1964, with 39 typhoons and tropical storms, and the least active was 2010, with 21 typhoons and tropical storms. There were 11 super typhoons in 1965 and only 5 in 2013. The 4-year period from 1964 to 1967 saw an average of 13.5 tropical storms, 15 typhoons and 7 super typhoons per year. The next most active period in the 1990s saw an average of 12 tropical storms, 15 typhoons and 5 super typhoons per year.²¹⁰

Storms forming over land also inflict casualties. The deadliest²¹¹ killed 1,300 people in the Manikganj District of Bangladesh in April 1989. A single lightning strike killed 21 people in a hut in present-day Zimbabwe in December 1975, and hail the size of cricket balls killed 246 people near Moradabad, India, in April 1888.²¹² A similar hailstorm hitting Sydney²¹³ on 14 April 1999 inflicted damages of \$1.7 billion (over \$4 billion today) but killed no one; the essential difference was wealth.

An analysis of hailpads in France from 1990 to 2010 by Hermida *et al.* (2013)²¹⁴ showed an upward trend in 154 (significant in 10%) and a downward trend in 177 (significant in 17%). Changnon and Changnon (2000)²¹⁵ assessed hail-day trends over a 100-yr period, 1896–1995, from carefully screened records of 66 stations across the United States, and found five types of 20-year fluctuation. They state:

One present in the Midwest had a peak in hail activity in 1916–35 followed by a general decline to 1976–95. Another distribution had a midcentury peak and was found at stations in three areas: the central high plains, northern Rockies, and East Coast. The third distribution peaked during 1956–75 and was found at stations in the northern and south-central high plains. The fourth temporal distribution showed a steady increase during the 100-yr period, peaking in 1976–95, and was found in an area from the Pacific Northwest to the central Rockies and southern plains. The fifth distribution found at stations in the eastern Gulf Coast had a maximum at the beginning of the century and declined thereafter. The 100-yr linear trends defined four regions across the United States with significant up trends in the high plains, central Rockies, and southeast, but with decreasing trends elsewhere in the nation. These up trends have occurred in areas where hail damage is greatest, and the trends matched well with those defined by crop-hail insurance losses and those found in studies of thunderstorm trends. The national average based on all station hail values formed a bell-shaped 100-yr distribution with hail occurrences peaking in midcentury.



Tornadoes in the U.S. have decreased in severity since recording began in 1954 (Fig. 18). NOAA states: “the increase in tornado reports over the last 54 years is almost entirely due to secular trends such as population increase, increased tornado awareness, and more robust and advanced reporting networks.”²¹⁶ Tornado activity increases with La Niña events, when the eastern Pacific Ocean is cool, and decreases during El Niño events.²¹⁷ NOAA reported 2011 as being the most costly year (over \$28 billion in damages) and the most deadly (with 551 fatalities) while 2016, the warmest year ever on record,²¹⁸ had the fewest tornadoes ever on record. There were 34 tornado-related fatalities in 2017.




Figure 18: Annual number of strong tornadoes across the United States, 1954-2014. Source: NCDC/NOAA²¹⁹

By reducing daytime and latitudinal temperature ranges, CO₂ may actually reduce the pressure differentials that drive extreme weather events. Rendering them less deadly requires a sound economy.

Floods take many lives and impact the health of survivors, often long after the event. The world’s deadliest flood occurred in 1332-3 when China’s Huang He (Yellow) River killed seven million people. The next deadliest floods occurred in 1887 and 1931 when the same river killed between a million and four million people.²²⁰ There were approximately 100,000 flood fatalities in England and the Netherlands in 1099, about 80,000 in the Netherlands when the “Great Storm” broke a dike in 1287, and another 10,000 from a similar incident in 1421. In 18th century Europe, late thaw ice jams blocked swollen rivers and burst dikes in the Netherlands.²²¹ The highest flood risk in Germany’s River Werra was in the 1700’s.²²² All but two of the 56 major floods affecting Florence since 1177 occurred before 1844.²²³ Flooding of the river Vltava in the Czech Republic decreased over the last century.²²⁴ All but two of Australia’s top ten worst floods²²⁵ occurred before 1975. During the 19th century, Brisbane had as many floods in 53 years as in the last 125 years. The highest flood was in 1841, followed closely by the prolonged and widespread flood of 1893 (Fig. 19).




Figure 19: Brisbane River heights in metres, 1840-2016. Source: Australian Bureau of Meteorology²²⁶

A global analysis of nearly 200 rivers revealed that flows over the last century were unchanged in the majority, increasing in 27 and decreasing in 31.²²⁷ This is also true for those rivers with observations stretching back much further in time.²²⁸ The IPCC's AR5²²⁹ agrees: "there continues to be a lack of evidence and thus low confidence regarding the sign of trend in the magnitude and/or frequency of floods on a global scale."

It is estimated that more than 59,000 flood fatalities occurred worldwide between 2005 and 2014,²³⁰ the vast majority in Asia, especially among women.²³¹ Haynes *et al.* (2016)²³² analysed 1,859 Australian flood fatalities from 1900 to 2015 and found a dramatic decline (Fig. 20) despite higher rainfall (Fig. 21). Ashley and Ashley (2008)²³³ found no statistically significant trend in fatalities over the period 1959-2005 in the United States (Fig. 22).




Figure 20: Australian flood fatality rate from 1900 to 2015. Source: Haynes *et al.* (2016)²³⁴




Figure 21: Annual mean rainfall across Australia, 1900-2017. Source: Australian Bureau of Meteorology²³⁵




Figure 22: Flood fatalities in the continental United States, 1959-2005. Black bars represent deaths due strictly to flooding for all event types in the study. Gray bars represent deaths due to tropical systems but not to flooding alone. Light gray bars represent deadly events. The dashed horizontal line represents yearly fatality median, and the nondashed horizontal line represents yearly fatality event median. The asterisk indicates that 2005 data are preliminary and do not include Hurricane Katrina fatalities from Louisiana. Source: Ashley and Ashley 2008²³³



Water security during flooding is important. Leptospirosis is a zoonotic disease acquired from infected animals, soil or water, especially flood water contaminated with rat urine. Globally, there are about a million severe cases and 60,000 deaths reported annually. Recent unprecedented outbreaks have been blamed on climate change, but the main reason is population growth plus poverty resulting in urban slums in developing countries.²³⁶

The Australian evidence indicates that the solution to flooding is not climate-mitigation but better infrastructure, warning systems, evacuation centres, rescue services etc. While flood casualties have been declining, health problems associated with dislocation and economic losses have been increasing due to population growth, especially along waterways. Flooding also has much more to do with altered land use and loss of flood-mitigating wetlands than with fossil fuel emissions. Bjorn Lomborg put it simply and starkly: “a dollar spent on flood management will reduce flooding 1,300 times better than a dollar spent on Kyoto.”²³⁷

Drought in the American west is often attributed to climate change, but accurate records of precipitation going back to 1895 indicate no significant trend in either the Southwest or Northwest Regions.²³⁸ Meko *et al.* (2007)²³⁹ demonstrated a robust relationship between tree-rings and observed stream flow in the Colorado River from 1905. Using these as proxies for stream flows back to 762CE, they found many severe droughts, the worst being in the 12th century (Fig. 23).




Figure 23: Colorado River stream flow, Observed: 1905-2005 & Reconstructed from Tree Rings: 762-2005
Source: Adapted from Meko et al, 2007.

The Sahel, covering 17 North West African countries, has a population of around 309 million, growing by about 3% per year. Devastating droughts there have been blamed on climate change.²⁴⁰ Rainfall over most of the Sahel declined at least 30% from the 1950s to the 1980s, but the 1950s were unusually wet, primarily because the Northern Hemisphere was warmer than the Southern Hemisphere; rainfall normalized as hemispheric temperatures equalized in the 1980s and 1990s.²⁴¹ A GFDL report,²⁴² updated 2 February 2018, concluded: “the drying from the 1950’s to the 1980’s is unlikely to be primarily forced



by increasing greenhouse gases in the atmosphere.” Climate models indicate that wet regions will become wetter and dry regions drier,²⁴³ but Sun *et al.* (2012)²⁴⁴ found that from 1940 to 2005, wet areas got drier while dry areas got wetter. Another study found no difference.²⁴⁵

As atmospheric CO₂ increases, leaf stomata density and water loss decrease;²⁴⁶ thus rendering plants more drought-tolerant.²⁴⁷ As CO₂ increased by 15% from 1982 to 2012, vegetation cover increased by 11% in arid areas.²⁴⁸ Konzmann *et al.* (2013)²⁴⁹ estimated that global irrigation demand will decline by about 17% by the 2080s, due to a combination of beneficial carbon dioxide effects on plants, shorter growing periods and regional precipitation increases. Regarding water availability, Wiltshire *et al.* (2013)²⁵⁰ estimate that the net global population at risk of high water stress will increase from 2.6 billion in 2000 to 4.1 billion in the 2080s due to population growth, but to 3.2 billion under the IPCC’s A1FI scenario (the one with the fastest warming), and to 2.9 billion when the benefits of CO₂ are factored in; the higher the CO₂ and the warmer the scenario, the greater the reduction in the population at risk of water stress.

Heatwaves are expected to increase in a warming world. Record-breaking heatwaves in Europe (2003²⁵¹, 2007²⁵²) Russia (2010²⁵³) the U.S. (2011²⁵⁴) Australia (2009²⁵⁵, 2013²⁵⁶) and globally²⁵⁷ have been linked to the human influence on climate, but this is far from certain. During the deadly 2003 European heatwave, for example, most of the planet was normal or cooler than normal at the time (Fig. 24). By comparison, 30% of the planet was 2 standard deviations (SDs) above normal (5% of it 3 SDs above) during the El Niño of 1998.





Figure 24: Global tropospheric temperature anomalies, June-August 2003. Source: Chase *et al.* (2006)²⁵⁸



Due to reduced soil moisture after another strong El Niño event, the Australian summer of 2016-17 was exceptionally warm, but only in the eastern states; Western Australia and the Northern Territory experienced an unusually cold summer and one of the wettest on record (Fig. 25). While Canberra reached 40°C on 9 February 2017, Perth reached just 17.4°C, its coldest February day on record. Twelve days later, Canberra had its coldest February morning on record! Heatwaves are weather events.




Figure 25: Mean Temperature Deciles for the 2016-17 Australian Summer. Source: Australian BOM

The worst heat wave in New South Wales since white settlement was in January 1896 when the maximum temperature was 102°F (38.9°C) or above for 24 days straight, 120°F (49.9°C) in the shade at Bourke, and 109°F (42.8°C) at midnight at Brewarrina.²⁵⁹ Hundreds died and the hospitals were crowded with victims.²⁶⁰ In the United States, a 10-day heatwave in the August of that year killed nearly 1,500 people,²⁶¹ mostly manual labourers in New York City.²⁶² Another very prolonged heatwave across the eastern states in 1901 killed 9,500 people, making it easily the most destructive heatwave in U.S. history.²⁶³ The 1936 North American heatwave during the Dust Bowl decade set record temperatures across 14 states, reaching 49°C in Steele, North Dakota.²⁶⁴ It caused catastrophic human suffering and killed at least 5,000 people. The summer-long heatwave across the Midwest of the USA in 1954 ranks as the hottest in 11 states from 1895 to 2009 (Fig. 26), reaching 117°F (47.2°C) on 14 July in East St Louis. Nancy Westcott (2011) analysed temperature records there from 1845 to 2009 and found a **reducing** trend over the 20th century. The UK had severe heatwaves in 1906, with temperatures reaching a September record of 35.6°C,²⁶⁵ and in 1911, with temperatures reaching an August record of 37.1°C.




Figure 26: Rank of the June–September 1954 heat wave based on National Climatic Data Center (NCDC) climate division temperature data for the years 1895–2009. Source: Westcott (2011)

Urban heat amplifies heatwaves. Large cities can be as much as 11°C warmer than the surrounding countryside.²⁶⁶ Torok (2001)²⁶⁷ found minimum temperatures in inner city Melbourne to be significantly higher than those in the suburbs. Edward Long (2010)²⁶⁸ compared NCDC rural and urban temperature datasets and found that the raw rural and urban data correlated very closely until 1965, when the urban data escalated rapidly. England's urban population jumped from 17% to over 70% during the 19th century. The global urban population was about 30% in 1950 and 50% in 2008, and is expected to reach 70%²⁶⁹ by 2050. Australia and the U.S. are now nearly 90% urban. Urban heat can be mitigated more effectively and economically than can greenhouse warming. Watts and Barrins (2014)²⁷⁰ describe how smart urban design can create cool refuges in cities. The greening of cities and cooling water features require adequate water resources, heatwaves often being associated with drought. We can learn many lessons from Cape Town.

Those at risk during heatwaves include the elderly, especially those living alone, people with heart or lung disease, pregnant or nursing mothers, babies and young children. Heat-tolerance is reduced by obesity²⁷¹ and increased by aerobic fitness.²⁷² Unlike cold spells, the mortality from heatwaves is reduced by acclimatisation.²⁷³ Whereas mortality remains higher for weeks after a cold spell, it drops rapidly and dramatically after a heatwave, indicating that most of those who die from short-term heat exposure were close to death.²⁷⁴ This “harvesting effect” has been demonstrated both in Europe²⁷⁵ and in the United States, so much so that “there is virtually no lasting impact of heat waves on mortality.”²⁷⁶

The excess mortality during the unusually hot European summer of 2003 was estimated at over 70,000.²⁷⁷ There were confounding variables: elevated atmospheric concentrations of ozone and particulate matter less than 10µm diameter (PM₁₀) accounted for 21-38% of deaths attributed to temperature



in the UK²⁷⁸ and for 33-50% of those in the Netherlands.²⁷⁹ In France, the death rate was also higher than that expected from temperature alone.²⁸⁰ Most of the 15,000 excess deaths occurred during the month-long August vacation during which there was a relative lack of care for the isolated and vulnerable, especially elderly females. The French Directorate General for Health then set up a National Heat Wave Plan which included a system for real-time surveillance of health data, compilation of scientific recommendations on the prevention and treatment of heat-related diseases, air-conditioning equipment for hospitals and retirement homes, emergency plans for retirement homes, city-scale censuses of the isolated and vulnerable, visits to those people during the alert periods, a warning system and repeated preventive message broadcasting by the media. When France experienced another severe heatwave in July 2006, Fouillet calculated nearly 4,400 fewer deaths than expected. Modern forecast and warning systems have been shown to be very effective in preventing heat-related deaths.²⁸¹ Compared to combating global warming, such adaptation measures are reliable, relatively cheap and rapidly effective in reducing mortality.

The 1954 U.S. heatwave was worse than the 1936 heatwave in intensity, duration and extent, but killed fewer than a fifth as many people,²⁸² largely because the home window air conditioners introduced after World War II became popular, as did fans, refrigerators, ice, cold drinks and cool clothing during the summer of 1954. While the global population has tripled since the 1920s, the global mortality from all weather-related natural disasters declined by 99%, from 242 per million to just 3 per million people per annum, thanks to improved economies and technologies.²⁸³ Instead of combating climate change to moderate heatwaves, the focus should be on evidence-based mitigation of urban heat and on proven adaptation measures.

Wildfires shaped the Australian landscape for 65 million years before indigenous people began using fire to do so about 45,000 years ago.²⁸⁴ Large parts of northern Australia burn every year or two during the dry season (winter and spring) after fuel accumulates during the summer monsoon season. As much as 35% of high-rainfall areas can burn in a typical year.²⁸⁵ Southern Australia experiences far fewer fires with a peak season in the summer and autumn, but they are much more deadly. More than 800 Australians have died in bushfires since 1850,²⁸⁶ the most in Victoria followed by NSW and Tasmania.²⁸⁷ Of the 173 Victorians killed on 'Black Saturday' in February 2009, 44% were under 12 or over 70 or unwell at the time.²⁸⁸

Unlike heatwaves, forest fires can have remote and protracted health consequences. The toxic and potentially carcinogenic smoke can impact health hundreds of kilometres away.²⁸⁹ Particulate matter smaller than 2.5 microns ($PM_{2.5}$) from wildfires and wood heaters is far more toxic to the respiratory system than equivalent concentrations of $PM_{2.5}$ from background urban sources.²⁹⁰ Johnson *et al.* (2012)²⁹¹ estimated that the smoke from wildfires kills over 300,000 people globally, mostly in Africa and Asia. On days when air pollution from surrounding bushfires is severe, cardiac arrests in Melbourne jump nearly 50%,²⁹² and Sydney's overall mortality rate increases by about 5%.²⁹³ Most affected are infants, the elderly and those with chronic heart and lung conditions,²⁹⁴ especially asthma and COPD.²⁹⁵ Wildfires in catchment areas can temporarily render the water unfit for human consumption²⁹⁶ and



reduce runoff by as much as 50% as trees regenerate over subsequent years or decades.²⁹⁷ Bushfires can also adversely impact the mental health of fire-fighters²⁹⁸ and survivors, especially those losing loved ones, livelihood and/or property, often for months and sometimes for years after the fire.²⁹⁹

The burning question, of course, is whether wildfire activity is increasing as a result of anthropogenic warming. Although the area burned from 1997 to 2011 decreased globally,³⁰⁰ increased forest fire frequency and/or duration of fire season has been observed in Africa,³⁰¹ South Africa,³⁰² Australia,³⁰³ Alaska,³⁰⁴ Canada,³⁰⁵ western USA,³⁰⁶ Russia³⁰⁷ and Spain.³⁰⁸ Factors influencing wildfires include the prevailing weather (hot dry winds), topography, fuel load (dry, oil-rich vegetation, altered ecosystems) and ignition source (natural or human). The human impact is also influenced by controlled burning, fire-breaks, clearing around dwellings, fire protection, warning systems, fire-fighting capabilities and demographics; most buildings affected by major bushfires in Australia are within 100m of bushland, and many back right onto it.³⁰⁹ Increasing atmospheric CO₂ levels are thought to increase wildfires by increasing fuel load via warmer and wetter growth conditions and by CO₂ fertilisation,³¹⁰ increasing combustibility via reduced precipitation and increased temperatures,³¹¹ and by increasing ignition via lightning strikes.³¹²

It is much more nuanced, however: increased atmospheric CO₂ benefits trees more than grasses, which dry out and burn faster. Ecosystem models are poor at simulating local variations in rainfall, tend to overestimate the growth response to precipitation, and don't include plant diversity, ground water, mineral composition of the soil, forest management, grazing, changes in cultivation practices and varieties, irrigation and disturbances such as storms and insect attacks, which together contribute the most variation in global greening.³¹³ Only a small minority of wildfires are ignited by lightning strikes, and this is thought to increase by perhaps 5-6% per degree (1°C) of warming.³¹⁴ Balch *et al.* (2017)³¹⁵ evaluated over 1.5 million government records of wildfires in the U.S. from 1992 to 2012 and found that humans accounted for 84% of them, that the human-caused fire season was three times longer than the lightning-caused fire season. Whereas those started by humans were dominant across most of the U.S. (5.1 million km²), those caused by lightning were dominant across only 0.7 million km², primarily the sparsely populated mountainous areas of the western United States. Many wildfires, such as California's Wine Country fires in 2017, are started by downed power lines in high winds.³¹⁶ A Royal Commission³¹⁷ into the 2009 Black Saturday bushfires in Victoria concluded:

Nine of the 15 fires the Commission examined were started as a direct or indirect result of human activity; five were associated with the failure of electricity assets, and the causes of four were thought to be suspicious. Broader data suggest that about one-third of bushfires in Victoria might be lit by people acting with mischievous or criminal intent. Although the proportion of fires that are caused by electricity infrastructure is low—possibly about 1.5 per cent of all ignitions in normal circumstances—on days of extreme fire danger the percentage of fires linked to electrical assets rises dramatically. Thus, electricity-caused fires are most likely to occur when the risk of a fire getting out of control and having deadly consequences is greatest.



Altered ecosystems can also have a profound impact; the introduction of African gamba grass³¹⁸ to northern Australia has greatly intensified wildfires there. Growing to 4m in height and competing strongly with native grasses, it can multiply the fuel load eightfold. New Zealand's Curran, Perry and Wyse³¹⁹ point out: "Plantations of highly flammable exotic species, such as pines and eucalypts, probably helped to fuel the recent catastrophic fires in Portugal and in Chile. In arid regions, such as parts of the U.S. southwest, the introduction of exotic grasses has transformed shrublands, as fires increase in severity."

An increased frequency, intensity or longer season of wildfires in some parts of the planet does not necessarily incriminate climate change. It may simply be that the population increased and so too did the number of firebugs, that expanding populations took aging electrical networks with them, that a recent tree-change brought people into closer contact with the bush, that misguided environmentalists prevented controlled burning, that flammable exotics invaded the landscape, and perhaps other reasons also. Focusing on CO₂ to prevent wildfires may prove to be as misguided as was focusing on eggs to prevent heart attacks. It will be a real tragedy if that focus diverts attention from proven preventive measures:

- Adequate and properly maintained firebreaks, including green firebreaks;
- Preventing or limiting the spread of highly flammable exotics;
- Maintaining safe electrical networks, placing them underground in high-risk areas;
- Regular planned/controlled burning at appropriate times/weather conditions;
- Timely effective communication to the public of fire risk, planned burns and smoke pollution;
- Making dwellings as fire-resistant and smoke-proof (sealed) as possible;
- Using effective portable air cleaners to reduce indoor smoke pollution, especially for asthmatics.


Such measures, of course, are costly and unaffordable by countries deprived of cheap energy, those African and Asian countries most at risk from wildfires and smoke inhalation. Rather than increasing the global mortality from wildfires, the use of fossil fuels might actually be reducing it in more ways than one.

Food, Famine, Climate and CO₂

Fifty years ago, in his book *The Population Bomb*, Paul Ehrlich predicted widespread famine with hundreds of millions starving to death in the 1970s. Instead, the death toll from famines declined even as the population grew (Fig. 27).



FIGURE 3.3 WORLD POPULATION GROWTH AND DEATH TOLL FROM GREAT FAMINES, 1900–2015



Note: Each great famine killed more than 100,000 people.
Source: US Census Bureau (2013a, 2013b); World Peace Foundation (2015).

Figure 27: Global population (billions) and mortality from great famines (millions)


While Ehrlich was wringing his hands, an agronomist from Minnesota, the late Norman Borlaug rolled up his sleeves and quietly launched a green revolution in Mexico, Pakistan, India and elsewhere. Mexico became a net wheat exporter in 1963 and the Philippines became a rice exporter in 1968. By 1970, India and Pakistan had more than doubled their wheat production, and “the man who saved a billion lives” was awarded the Nobel Peace Prize for his work. Thanks to fossil fuels and Borlaug’s disease-resistant high-yield hybrids and fertilisers, global food production increased by 2.3% pa from 1961 to 2005, which was faster than the 1.7% pa population increase. Food consumption increased from 2280 kcal/d to 2800 kcal/d per person, and a strong correlation was observed between agricultural growth rates and GDP; the only region where food production per capita has not increased is sub-Saharan Africa.³²⁰ Africa’s food production was predicted in 2008 to decline by as much as 50% by 2020 due to climate change.³²¹ The major threat, however, is not climate change but civil war and the armyworm, a native maize-munching moth of the Americas which turned up in Nigeria early in 2016 and spread rapidly across the continent. The American industry was saved by a bacterium gene which scientists inserted in Bt maize, making it toxic to the fall armyworm but not to humans. South Africa, the only African country to approve Bt maize, had a record crop in 2017.³²²

In 2013, the IPCC’s AR5³²³ stated: “For the major crops (wheat, rice, and maize) in tropical and temperate regions, climate change without adaptation is projected to negatively impact aggregate production for local temperature increases of 2°C or more above late-20th-century levels.” In *Nature Climate Change*, Asseng *et al.* (2015)³²⁴ warned: “warming is already slowing yield gains at a majority of wheat-growing locations. Global wheat production is estimated to fall by 6% for each °C of further temperature increase.” This prompted a media report³²⁵ that India “produced one million metric tons more wheat in 2013 than it did in 2014.” But this was due to drought. After global temperatures climbed to record levels, *The Times of India* (17 August 2017)³²⁶ reported a record food-grain production of 275.68 million tonnes, up



4% on the previous record, a five-fold increase since 1951 and a four-fold yield per hectare. The global wheat production reached record levels due to increases in India, Russia and the U.S. (Fig. 28). In March 2018, *The Guardian*³²⁷ reported on durum varieties that can withstand constant 40°C temperatures.

Fig 4--Global wheat production and use



Graph from Food Security Porta, with latest USDA figure estimated added. (joannenova.com.au)

Figure 28: Global wheat production this century (thousand tons pa). Source: World Grain, November 2016³²⁸


Global warming extended the arable area, growth rate and growing season for food crops.³²⁹ From 1980 to 2003, global food production increased by 62.8% as atmospheric CO₂ increased by 11.2%, fertiliser use by 27.5% and land use barely at all.³³⁰ Numerous studies of CO₂ enrichment in chambers (e.g. greenhouses) have demonstrated dramatically improved crop yields.^{331, 332} Ainsworth-Long (2005)³³³ performed a meta-analysis of 124 papers on 40 species tested at 12 sites, 7 in the USA, 3 in Europe, 2 in New Zealand and Japan, using free-air CO₂ enrichment (FACE). The actual increases achieved (above the ambient CO₂ level at the time of the study) varied from 30.5% to 68% with a median of 50-55% (550ppm) and an average of 49.2%. They found that trees benefited the most, more than anticipated from chamber studies, while average crop yields increased by 17%, less than expected from chamber studies. The overall trend level for wheat was about 15%. Sorghum yields increased by as much as 28% under dry conditions, due to reduced water loss from fewer transpiration stomata. Light-saturated CO₂ uptake was increased by 19% at temperatures under 25°C and by 30% at higher temperatures, indicating much better heat-tolerance under elevated CO₂. Elevated CO₂ also reduced the toxic effects of ozone. Two FACE facilities using CO₂ at up to 200ppm above ambient levels produced a 5-7% increase in rice yield and 8% increase in wheat yield, which would result in an extra 59 million metric tons a year globally, enough to feed an extra 550 million mouths at the average per capita consumption. Reducing atmospheric CO₂ to preindustrial levels, advocated by some climate activists, would put billions of lives at risk of starvation and thus constitute a crime against humanity. See our White Paper: *The Climate Surprise – Why CO₂ is Good for the Earth*.³³⁴



Moreover, CO₂ fertilisation has been shown to significantly increase the flavonoid content of wheat³³⁵ and strawberries.³³⁶ Wang *et al.* (2003) found that increasing the CO₂ concentration by 300ppm and 600ppm increased strawberry flavonoids by 55% and 112%, ascorbic acid (vitamin C) by 10% and 13% and glutathione (another antioxidant) by 3% and 171% respectively. Tomatoes grown in enriched CO₂ are higher in vitamins A³³⁷ and C.³³⁸ Idso *et al.* (2002)³³⁹ found that CO₂-enriched oranges were 4% heavier, 74% more in number and 5% higher in vitamin C. Soy beans grown from seedlings in CO₂ at 700ppm (compared to 400ppm) had an isoflavone content 8% higher when grown at the usual mean temperature of 18°C, 104% higher when grown at 23°C and 101% higher at 28°C.³⁴⁰ When drought was added to heat stress, the isoflavone content was 38-186% higher in plants exposed to 700ppm. Kim *et al.* (2005) likewise found a 72% increase in soy isoflavones grown at 650ppm (vs. 360ppm), and a 96% increase in total plant biomass. Broccoli grown in 65%-enriched CO₂ produced heads 7% heavier and containing 37% more glucosinolates.³⁴¹ These enhance flavour and are thought to help prevent cancer.³⁴² Similar results were found with Chinese kale.³⁴³ Growing spinach at 800ppm increased the fresh weight by 67%, the soluble protein concentration by about 52% and vitamin C by 21%.³⁴⁴ Gwynn-Jones *et al.* (2012)³⁴⁵ found that quercetin glycosides and various other antioxidants were significantly higher in several types of berry consumed by humans and other animals at Northern Latitudes when grown at 600ppm. It is very likely that CO₂ has been quietly improving food quantity and quality.

Global food security depends not only on production but also on distribution and transport, which invariably involves the use of fossil fuels. Famine and starvation in the developing world is usually the product of politics plus natural disaster, sometimes aggravated by climate change but more often by misguided climate action. The most notable example is the diversion of good food into biofuel. Goklany (2011)³⁴⁶ estimated that such activities in 2008 helped to push 130-155 million people into absolute poverty, hunger and starvation, and may have led to 190,000 premature deaths in 2010 alone. Australia is a major wheat exporter, but a 2007 CSIRO report warned that “If all of the ethanol capacity that is currently proposed was to be fulfilled by existing crops (principally wheat and sugar), or if a national E10 target were to be met (eg. by 5.5 Mt of wheat as the feedstock), it could force the import of wheat in drought years.” Despite a severe drought in 2012 in the U.S., 40% of its corn crop³⁴⁷ went into ethanol. The real irony is that “Once estimates from the literature for process emissions and displacement effects including land-use change are considered, the conclusion is that U.S. biofuel use to date is associated with a net increase rather than a net decrease in CO₂ emissions.”³⁴⁸ It is worse than useless! Such folly persists only because of politics and taxpayer subsidies to big business.

Seafood plays an important role in food security, supplying about 10% of world human calorific intake.³⁴⁹ The omega-3 fatty acids (ω-3 FAs) produced by plankton at the bottom of the food chain probably reduce³⁵⁰ cardiovascular morbidity and mortality and may benefit some metabolic, inflammatory, neurological, neuropsychiatric and eye disorders.³⁵¹ The UK Scientific Advisory Committee on Nutrition guidelines (2004) recommends at least two portions of fish a week. It is thought that increasing CO₂ will impact seafood via higher sea surface temperatures, ocean ‘acidification’ (actually slightly reduced



alkalinity), altered precipitation/runoff and sea level rise. Crustaceans (crabs, lobsters, shrimps and krill) appear to benefit from more dissolved CO₂ even at seven times today's atmospheric level.³⁵² Relying on extreme emission/temperature scenarios, Cheung *et al.* (2009)³⁵³ report:

Here, we project changes in global catch potential for 1066 species of exploited marine fish and invertebrates from 2005 to 2055 under climate change scenarios. We show that climate change may lead to large-scale redistribution of global catch potential, with an average of 30–70% increase in high-latitude regions and a drop of up to 40% in the tropics. Moreover, maximum catch potential declines considerably in the southward margins of semienclosed seas while it increases in poleward tips of continental shelf margins. Such changes are most apparent in the Pacific Ocean. ... Many highly impacted regions, particularly those in the tropics, are socioeconomically vulnerable to these changes. Thus, our results indicate the need to develop adaptation policy that could minimize climate change impacts through fisheries.

Increasing atmospheric CO₂ is having uncertain positive and negative impacts on seafood supplies. One thing is certain: “global marine fisheries are underperforming economically because of overfishing, pollution and habitat degradation.”³⁵⁴ We need to focus on real problems, particularly plastic pollution of the oceans. For a comprehensive review, see *Ocean Acidification* by Patrick Moore in our White Paper: ***The Climate Surprise – Why CO₂ is Good for the Earth.***³⁵⁵

Social Strife and Mental Health

Climate change is predicted to cause major social dislocations and mental health problems. In a forecast prepared for the Pentagon in 2003, Schwartz and Randall³⁵⁶ predicted (among many others):

In 2007, a particularly severe storm causes the ocean to break through levees in the Netherlands making a few key coastal cities such as The Hague unlivable [sic]. ... Melting along the Himalayan glaciers accelerates, causing some Tibetan people to relocate. . . .

2010: Border skirmishes and conflict in Bangladesh, India, and China, as mass migration occurs toward Burma

2012: Severe drought and cold push Scandinavian populations southward, push back from EU

2012: Flood of refugees to southeast U.S. and Mexico from Caribbean islands

2015: Conflict within the EU over food and water supply leads to skirmishes and strained diplomatic relations

Needless to say, none of these prophesies were fulfilled. Owain and Maslin (2018)³⁵⁷ found “that climate variations played little or no part in the causation of conflict and displacement of people in East Africa over the last 50 years. Instead, we suggest rapid population growth, low or falling economic growth and political instability during the post-colonial transition were the more important controls.” In his



2004 bestseller, *The Weather Makers*, Tim Flannery predicted:

When we consider the fate of the planet as a whole, we must be under no illusions as to what is at stake . . . the fate of hundreds of thousands of species, and most probably billions of people. (p. 170) It seems inevitable that these [Pacific Island] nations will be destroyed by climate change during the course of this century. (p. 287)

Besides many more wild exaggerations, this book contained numerous contradictory statements and factual errors.³⁵⁸ In 2006, Al Gore's docudrama, *An Inconvenient Truth*,³⁵⁹ emotively depicted polar bears drowning, oceans rising 20 feet, dreadful epidemics and more – all happening very soon unless we 'save the planet'. After being shown in Australian schools, a 2007 survey of 600 children aged 10 to 14 revealed that 44% were nervous about the future impact of climate change and, "a quarter honestly believe the world will come to an end before they get older."³⁶⁰ A subsequent survey of 200 schoolchildren in New South Wales likewise found that fears over climate change were producing feelings of powerlessness and despair: "Many children thought they would not survive to adulthood."³⁶¹ What does this do to career aspirations! Do predictions of perpetual drought cause farmers to abandon hope? The climate of fear campaign has produced more anxiety and despair than has climate change itself.

Climate change catastrophism has also polarised society. Merchants of fear attack Merchants of Doubt.³⁶² Sceptics of *catastrophic* climate change are suspected of big-oil funding but labelled paranoid if they suspect scientists of being biased by big-government funding. Robust scientific debate has been replaced by a claimed 97% scientific consensus,³⁶³ where there is no consensus on **how much** anthropogenic warming has occurred or will occur,³⁶⁴ especially when carefully analysed.³⁶⁵

Society is also becoming increasingly litigious over climate change: the Commonwealth Bank of Australia³⁶⁶ and even financial advisors³⁶⁷ for failing to disclose climate change risk in reports; the UK government³⁶⁸ and ministers³⁶⁹ for failure to fight climate change; the Dutch³⁷⁰ and U.S. governments;³⁷¹ Californian cities and counties claiming damages from rising seas, storms and wildfires against oil companies³⁷² and receiving counter-claims.³⁷³ As early as 2002, Tuvalu³⁷⁴ threatened to take legal action against the worst emitters of greenhouse gases, the United States and Australia, because their island would soon be swallowed by rising seas and their 11,500 people would become the world's first climate refugees. Sea level around Tuvalu actually fell during the latter half of the 20th century.³⁷⁵ For more information on sea level, see our White Paper: ***A Primer on Carbon Dioxide and Climate***.³⁷⁶

Energy Sources and Health

Existing energy sources include:

- Fossil Fuels – coal, oil and gas
- Biomass (wood, crop waste and dung) and Biofuel (ethanol and biodiesel)
- Wind and solar (photovoltaic and thermal solar)
- Hydropower including pumped hydro
- Nuclear power



Coal power has saved countless lives from drudgery, and taken many lives via the inhalation of soot. In a letter to the Financial Times (November 12, 2015), Lord Stern claimed that 7 million deaths each year were caused by the air pollution from fossil fuels, mainly coal. On 25 November, Bjorn Lomborg responded:

The facts of the matter, as established by the World Health Organisation, are that the majority of air pollution deaths comes from indoor pollution, and about 85 per cent of these deaths are caused by biomass burning. This translates to 3.2m of the 7m deaths caused by indoor biomass burning. Moreover, a large part of the outdoor air pollution stems from non-fossil fuels. The most recent Global Burden of Disease estimates that 12 per cent of all outdoor air pollution comes from indoor air pollution, causing an extra 373,000 premature deaths. The most recent study from Nature estimates that just the indoor air pollution from households in China and India spilling into the open air causes 760,000 outdoor air pollution deaths. The Nature study shows that 600,000 outdoor air pollution deaths are caused by natural sources (mostly airborne desert dust). Another 660,000 deaths are caused by agriculture, mostly from release of ammonia, forming ammonium sulphate and nitrate. Finally, almost 200,000 additional deaths come from large biomass burning (forest burning such as we recently saw in Indonesia).

Power generation, traffic and industry, which are mostly fossil fuel-driven and likely what Lord Stern was thinking about, in total cause 854,000 air pollution deaths. Added to the 560,000 deaths from indoor air pollution caused by coal, this constitutes only 20 per cent of total air pollution deaths, and hence is a far cry from Lord Stern's claim of a majority.

This matters for two reasons. First, it is disingenuous to link the world's biggest environmental problem of air pollution to fossil fuels and indirectly climate. It is a question of poverty (most indoor air pollution) and technology (scrubbing pollution from smokestacks and catalytic converters) — not about global warming and CO₂. Second, costs and benefits matter. Tackling indoor air pollution turns out to be very cheap and effective, whereas tackling outdoor air pollution is more expensive and less effective.

According to the World Health Organisation (WHO),³⁷⁷ over 3 billion people still cook and heat their homes by burning biomass, and over 1.2 billion people have no access to electricity. Africa has 17% of the world's people but only 3% of its electricity.³⁷⁸ By replacing biomass for cooking and heating, and supplying 39% of the world's electricity, coal prevents millions of deaths every year. By helping to lift people out of poverty, it saves millions more. High-efficiency low-emission (HELE) power plants³⁷⁹ use scrubbers and precipitators to remove 99% of the fly ash, 97% of the sulphur dioxide and up to 90% of the nitrogen oxides. New ultra-clean coal (UCC) technologies can remove 99.75% of particulates and almost all the sulphur. Coal power in the U.S. is 17 times safer than in India and China (Table 1). India has huge coal reserves but mostly low-quality with high ash content,³⁸⁰ so needs to import high-quality coal and/or build HELE plants.

There are hazards associated with the mining and transport of coal. Wikipedia lists 50 mining disasters³⁸¹ since 1885, mostly underground cave-ins and explosions. The worst year on record was



1907 when over 3,000 U.S. miners died.³⁸² Coal worker pneumoconiosis (CWP), commonly known as ‘black lung’, took about 25,000 lives globally in 2013.³⁸³ After being eliminated in Queensland, Australia in the 1970s and in New South Wales a little later (Fig. 29), it recently re-emerged,³⁸⁴ being confirmed in 18 Qld miners and one in NSW. The U.S. has been less successful in preventing CWP, probably due to the higher quartz content³⁸⁵ in mine dust, especially in the central Appalachian region where one in ten miners are affected.³⁸⁶ Modern open cut mining is much safer but not without accidental incidents. Particles less than 10 microns in diameter (PM10) pollute the environment. Spraying a bio-degradable binding polymer onto the surface has been shown to reduce coal dust lost to the environment by 75%. Queensland includes this in its extensive management controls.³⁸⁷




Figure 29: Prevalence of pneumoconiosis among U.S. underground coal miners and NSW coal workers.
Source: Joy, Colinet and Landen: CDC³⁸⁸

Oil and its distillates (petroleum, kerosene and diesel) revolutionised 20th century transport even more than did coal in the 19th century. Lead (added to petrol as anti-knock) contaminated the air and soil along busy roadways and smog plagued large cities until unleaded petrol and expensive three-way catalytic converters³⁸⁹ were mandated in developed countries during the 1980s.

After Kyoto, the EU decided in 2001³⁹⁰ to promote diesel to reduce CO₂ emissions. By 2016, most cars made for the European market were diesel. But diesel exhaust contains more nitrogen oxides and relatively high levels of particulates (soot) which catalytic converters can’t remove. Diesel particulate filters (DPF) were mandated in the United States in 2007, but they tend to clog up during short trips around town, so periodic regeneration requiring more fuel is used to remove the accumulated carbon



from the DPF. This led to computer tampering during testing. An honest study showed the latest “Euro 6 Standard” vehicles miss their pollution targets by a whopping 400%.³⁹¹ In a misguided attempt to reduce a non-pollutant (CO₂), Europe massively increased real pollution! After successive Indian governments subsidised diesel to assist farmers, the number of diesel cars increased ten-fold between 2000 and 2013. Delhi now has the world’s worst air-pollution, killing up to 50,000 people a year according to the WHO.³⁹² Biodiesel is no less polluting. Cities could be made cleaner, cooler and quieter by replacing combustion engines with electric motors. While electric vehicles today are mostly coal-powered, they will eventually be mostly nuclear-powered.

Gas (methane), which produces almost no pollutants and less CO₂ per unit energy (Table 1), is fast replacing coal in the United States. Fracking³⁹³ for shale gas has a good safety record and opposition is waning.³⁹⁴ Being far more flammable than other fossil fuels, its storage and distribution requires stringent safety standards. The CBS News³⁹⁵ reported: “In 2009, there were 158 natural-gas distribution pipeline incidents reported to PHMSA that killed 10 people, injured 50 and caused almost \$32 million in property damage.”

Table 1: Pounds of CO₂ emitted per million British thermal units (Btu) of energy for various fuels.

Source: U.S. Energy Information Administration³⁹⁶

Fuel Source	Lbs. CO ₂
Coal (anthracite)	228.6
Coal (lignite)	215.4
Coal (bituminous)	205.7
Diesel fuel and heating oil	161.3
Gasoline (without ethanol)	157.2
Propane	139.0
Natural gas	117.0

Biomass accounts for about 10% of global energy consumption³⁹⁷ and around 90% in South Asia and sub-Saharan Africa (except South Africa).³⁹⁸ It is unquestionably the most air-polluting and lethal of all energy sources. The WHO³⁹⁹ states: “4.3 million people a year die prematurely from illness attributable to the household air pollution caused by the inefficient use of solid fuels (2012 data) for cooking. Among these deaths, 12% are due to pneumonia, 34% from stroke, 26% from ischaemic heart disease, 22% from chronic obstructive pulmonary disease (COPD), and 6% from lung cancer.” The WHO⁴⁰⁰ estimates that polluted air impacts 90% of people, kills around 7 million annually and accounts for 36% of lung cancer deaths, 34% of stroke deaths, 27% of heart disease deaths and 11.6% of all deaths globally.

Ambient (outdoor) air pollution comes from dust, forest fires and burning biomass as well as from fossil



fuels. Lelieveld *et al.* (2015)⁴⁰¹ attributed 90% of the ambient air pollution in South Asian megacities to the burning of biomass in homes. Mikko Paunio,⁴⁰² an epidemiologist at the University of Helsinki, recently stated:

Thus perhaps around six million deaths globally are attributable to domestic combustion of solid (bio)fuels. However, despite these appalling statistics, the development community has focused its efforts on mitigating global warming instead. Some . . . have even encouraged the burning of crop residues in homes. The effect of this headlong rush to 'save the climate' has horrifying implications for human health.

Paunio also points out that burning wood pellets in Europe is not only more polluting than coal but also increasing CO₂ emissions.⁴⁰³ The same applies to most biofuel, as discussed under food and famine.

A large prospective study from 2000 to 2009 across the United States indicated that fine particulate matter (≤ 2.5 microns or PM_{2.5}) levels over $10\mu\text{g}/\text{m}^3$ increased total mortality by 3%, CVD mortality by 10% and respiratory mortality in never smokers by 27%.⁴⁰⁴ The most air-polluted region of the planet, with an annual mean ambient PM 2.5 of greater than $26\mu\text{g}/\text{m}^3$, stretches band-like from northwest Africa across Arabia and northern India to eastern China; countries with the highest CO₂ emissions per capita, Australia, Canada and United States, have among the lowest actual air pollution (i.e. PM_{2.5} $\leq 10\mu\text{g}/\text{m}^3$); only Siberia, Scandinavia, Scotland and Ireland have less, and the only areas in Australia with a PM_{2.5} greater than $10\mu\text{g}/\text{m}^3$ are in the wildfire-affected north and the dry undeveloped centre.⁴⁰⁵ The answer to life-threatening air pollution is development, and that requires the intelligent use of fossil fuels.

Solar power includes both photovoltaic (PV) and thermal solar plants, some of which can store heat for use after sundown. There are hazards associated with PV in the manufacture, rooftop installation, cleaning, servicing and removal/replacement.⁴⁰⁶ Potentially toxic materials used in PV manufacture include silicon, gallium arsenide and cadmium,⁴⁰⁷ cadmium telluride providing the best energy return on energy invested (EROI = 32:1).⁴⁰⁸ The PV solar industry has also become one of the leading emitters⁴⁰⁹ of hexafluoroethane, nitrogen trifluoride and sulphur hexafluoride, potent and potentially toxic greenhouse gases.⁴¹⁰

In the U.S. construction industry, falls are the leading cause of fatalities, 269 from ladders and roofs in 2002. Brian Wang (2008)⁴¹¹ estimated 100-150 fatal falls from solar panel roof installations annually. As PV solar is only about 15% efficient, most insolation on black solar panels is converted to heat, thus contributing to urban heat. Lasting less than 30 years⁴¹² and containing lead, chromium and cadmium, they create 300 times more toxic waste per unit of energy produced than do nuclear power plants; a solar waste crisis looms.⁴¹³ PV solar panels also require the most resources per unit of energy produced (Fig. 30).




Figure 30: Material resources (tonnes) per unit energy produced (TWh). Source: Desai and Nelson (2017).⁴¹⁴

Wind power impacts human health in the manufacture and installation of turbines, in the visual amenity, noise, economic and agricultural impacts; many European farms are growing turbines instead of turnips! Most fatalities associated with wind (Table 2) occur during the transportation and installation of turbines. With nacelles the size of buses and blades 50m long, reaching heights of 150m and sweeping an area of 1.9 acres with each rotation, wind turbines blight the landscape. Sunlight flickering through rotating blades can be annoying, and at 3Hz can trigger an epileptic seizure in susceptible subjects.⁴¹⁵ While one acclaimed U.S. study found that wind turbines had no impact on real estate values,⁴¹⁶ a UK study⁴¹⁷ showed wind turbine visibility reduced house prices substantially.⁴¹⁸

Studies and reports on wind turbine noise are many, varied and often biased.⁴¹⁹ A comprehensive review by Schmidt and Klokke (2014)⁴²⁰ found: “Wind turbines emit noise, including low-frequency noise, which decreases incrementally with increases in distance from the wind turbines. Likewise, evidence of a dose-response relationship between wind turbine noise linked to noise annoyance, sleep disturbance and possibly even psychological distress was present in the literature. Currently, there is no further existing statistically-significant evidence indicating any association between wind turbine noise exposure and tinnitus, hearing loss, vertigo or headache.” Infrasound (<20Hz) at 1000m from large wind turbines (>2MW) or 500m from smaller ones was very similar to that 8km inland from the coast (56-58dB), and less than that at 350m from a gas-fired power plant, 25m from the waterline at



the beach or 70m from major roads (74-76dB). Rather than aggravating Meniere's disease, infrasound pulses at 6Hz can be therapeutic.⁴²² Infrasound in the 5–8 Hz range can nevertheless cause a rattling of doors and windows which is audible and can be annoying to those living in close proximity to wind turbines.⁴²³ Schmidt and Klokke, noting the inability of cross-sectional studies to determine a clear causal relationship, state: "It is therefore not known with certainty if the association between wind turbine exposure and health-related outcomes is caused by sound exposure, visual disturbance, economic aspects or something else."

Tang *et al.* (2017)⁴²⁴ found that wind turbines in northern China affected the soil temperature and moisture of nearby farms, thereby reducing gross summer production by 8.9% and annual net production by 4%. Decommissioning⁴²⁵ of Europe's earliest offshore wind turbines, after about 25 years of operation, reveal an EROI of about 1:1 or less. The major problem with wind and solar is its intermittency and unreliability. Nearly half of South Australia's electricity came from wind turbines when strong wind forced their shut down in September 2016, resulting in a state-wide blackout.⁴²⁶ South Australia has the highest electricity price and unemployment rate in Australia. California, Denmark and Germany likewise have the world's highest⁴²⁷ use of renewables and electricity prices. Expensive power increases fuel poverty and cold-related mortality.

Hydropower is the most reliable renewable energy, provided the rain comes. MacLean *et al.* (April 2018) reported in The Washington Post: "Low water levels at these big dams, such as Ghana's Akosombo Dam,⁴²⁸ built under the guidance of the country's first president, Kwame Nkrumah, cause frequent load shedding and blackouts. In November 2017, Tanzania experienced a nationwide blackout⁴²⁹ and has faced recurrent blackouts for a decade because of low water levels from drought."⁴³⁰ Pumped hydro allows surplus energy to be stored while the wind blows and the sun shines. It is vital that dams and reservoirs are well designed and built only on suitable sites. When the Banqiao dam on the Ru River in China burst in 1975, more than 170,000 people perished. A 1972 dam failure in Buffalo Creek, West Virginia took 125 lives, injured over 1,100 and left nearly 5,000 people homeless. In 1972, a dam near Rapid City, South Dakota, failed, which flooded the entire downtown during the night. This event killed 237 and injured 2,932 people.

Nuclear power has been on the nose since a tsunami hit Fukushima on 11 March 2011, even though no one died from radiation.^{431, 432} Nuclear has by far the lowest fatality footprint, especially in responsible countries such as the U.S. (Table 2). Given that there have been only 50 confirmed deaths⁴³³ from nuclear power plants, including Chernobyl, the estimate of 90 per PWh in Table 2, prepared by Conca,⁴³⁴ who relied on Brian Wang (2008) and some questionable assumptions, is a gross exaggeration. According to Conca, wind and nuclear have the smallest carbon footprint (~15 g emitted per kWhr). Unlike early wind turbines, nuclear power has an EROI of 70:1⁴³⁵ and requires far fewer resources (Fig. 30). Laser enrichment⁴³⁶ of radioactive waste and fusion breeding⁴³⁷ might soon revolutionise the nuclear industry, making it more efficient and safer than ever.



Table 2: Mortality rate per unit of power produced by energy sources (2012)

The numbers combine actual deaths and epidemiological estimates, and are rounded to two significant figures

Energy Source	Deaths/PetaWh	Percentage of energy type
Coal – global average	100,000	41% of global electricity
– China	170,000	75% of China’s electricity
– U.S.	10,000	32% of U.S. electricity
Oil	36,000	33% of global energy – 8% of electricity
Natural Gas	4,000	22% of global electricity
Biofuel and Biomass [#]	24,000	21% of global energy
Solar - rooftop	440	<1% of global energy
Wind	150	3% of global electricity
Hydro – global	1,400	16% of global electricity
– U.S.	5	6% of U.S. electricity
Nuclear – global	90	17% of global electricity
– U.S.	0.1	19% of U.S. electricity

[#]Biofuel greatly dilutes the Biomass fatality rate.

Source: James Conca in Forbes¹

For optimal health, every country should be free to choose the most affordable and reliable energy sources. Those using coal should be encouraged and perhaps assisted to make it as clean as possible. Most will probably transition to gas and eventually to nuclear energy. Biomass should be phased out, as should all subsidies, and intermittent renewables must remain minor players. The underlying principle should be the conservation and best use of resources.

Conclusion

Warmth is good for human health and prosperity. Fossil fuels have played a vital role in providing the wealth essential for health and environmental protection. They have also boosted atmospheric CO₂ and added a little warmth, both being hitherto beneficial overall for plants and people. The ingenuity of Homo sapiens at adapting to climate has permitted people to populate almost the entire globe. No other animal has managed to thrive in the freezing Arctic and steamy tropics. If we stick to doing what we do best – adaptation – we will not only survive global warming, we will continue to thrive. We must be prepared not only for global warming, but also for global cooling, which will occur as our present warm Holocene draws to its inevitable end.

Human health and that of the planet depends on balancing productivity and development with conservation and environmental protection. Only developed countries with people lifted out of poverty can afford to produce clean energy, protect the environment, put power underground, construct buildings with 5-star energy ratings and use efficient lighting/appliances to minimise energy and water use, provide adequate safe water supplies and effective public health measures to control communicable diseases. It is vital that governments focus on real pollutants, not imagined ones, and that they avoid using climate



change as a scapegoat for failure to implement sound public health policies and proven preventive measures. Misguided climate action can be worse than unmitigated climate change. When women (in particular) are educated and agriculture is mechanised, as in the developed world, fertility rates drop and population growth slows, thus reducing social and environmental pressures, conflict and emigration.

Urban design can be improved to reduce urban heat, and to encourage health-promoting walking and cycling. Smoggy cities could also encourage a switch to electric vehicles, but not with generous taxpayer-funded subsidies. Energy costs need to be kept as low as possible, especially in cold climates, so that poor people can afford to keep warm in winter. For every death from heat, there are twenty from cold. Fossil fuels, including clean-coal will continue to have an important role to play in advancing civilisation and human health over the 21st century. Our focus should be on conservation and health-promoting activities rather than on CO₂ and climate change. Unmitigated warming this century is likely to be less than 1°C and thus more beneficial than harmful for humanity and perhaps for the planet.

We concur with the latest IPCC Summary for Policymakers:⁴³⁸ “The most effective vulnerability reduction measures for health in the near term are programs that implement and improve basic public health measures such as provision of clean water and sanitation, secure essential health care including vaccination and child health services, increase capacity for disaster preparedness and response, and alleviate poverty (very high confidence).”

References

1. Dansgaard, W., White, J. W. C. and Johnsen, S. J. 1989: The abrupt termination of the Younger Dryas climate event. *Nature* 339: 532-533.
2. Lamb, H. H. 1988: *Weather, Climate and Human Affairs: A Book of Essays and Other Papers*. London and New York: Routledge.
3. Cuffey, K. M. and Marshall, S. J. 2000: Substantial contribution to sea level rise during the last interglacial from the Greenland ice sheet. *Nature* 404: 591-594
4. McBean, G. *et al.* 2005: *Arctic Climate: Past and Present*, Chapter 2, p. 51. http://www.acia.uaf.edu/PDFs/ACIA_Science_Chapters_Final/ACIA_Ch02_Final.pdf 2005; accessed June 12, 2018.
5. Briner, J. P. *et al.* 2006: A multi-proxy lacustrine record of Holocene climate change on northeastern Baffin Island. *Quaternary Research* 65: 431-442
6. Rosen, W. 2007: *Justinian's Flea: Plague, Empire, and the Birth of Europe*. Viking Adult, p. 3, ISBN 978-0-670-03855-8.
7. Lamb, H. H. 1965: The early Medieval Warm Epoch and its sequel. *Palaeogeology, Palaeoclimatology, Palaeoecology* 1:13-37.
8. Patterson, W. P., Dietrich, K. A., Holmden, C. and Andrews, J. T. 2010: Two millennia of North Atlantic seasonality and implications for Norse colonies. *Proc. National Academy of Science* www.pnas.org/cgi/doi/10.1073/pnas.0902522107 accessed June 12, 2018.
9. Wikipedia: Great Famine of 1315–17. https://en.wikipedia.org/wiki/Great_Famine_of_1315%E2%80%9317 May 22, 2018; accessed June 12, 2018.
10. Wikipedia: Great Famine of 1315–17. https://en.wikipedia.org/wiki/Great_Famine_of_1315%E2%80%9317 May 22, 2018; accessed June 12, 2018.



11. Austin Alchon, S. 2003: *A pest in the new land: new epidemics in a global perspective*. University of New Mexico Press. p. 21. ISBN 0-8263-2871-7. See <https://books.google.com/books?id=YiHHnV08ebkC&pg=PA21&dq#v=onepage&q&f=false> Accessed June 12, 2018.
12. Gordon, D. 1976: *Health, Sickness and Society*. University of Queensland Press, St. Lucia, p.185.
13. Bull, G.M. and Morton, J. 1975: Relationships of temperature with death rates from all causes and from certain respiratory and arteriosclerotic diseases in different age groups. *Age and Ageing* 4: 232–246.
14. Wikipedia: https://en.wikipedia.org/wiki/Population_growth May 30, 2018; accessed June 12, 2018.
15. Fernandez-Raga, M., Tomas, C. and Fraile, R. 2010: Human mortality seasonality in Castile-Leon, Spain, between 1980 and 1998: the influence of temperature, pressure and humidity. *International J. of Biometeorology* 54: 379–392.
16. Moore, T.G. 1998: *Climate of Fear*, Cato Institute: Washington DC p. 83.
17. Davis, R.E., Knappenberger, P.C., Michaels, P.J., and Novicoff, W.M. 2003: Changing heat-related mortality in the United States. *Environmental Health Perspectives* 111: 1712–1718
18. Davis, R.E., Knappenberger, P.C., Michaels, P.J. and Novicoff, W.M. 2004: Seasonality of climate-human mortality relationships in U.S. cities and impacts of climate change. *Climate Research* 26: 61–76.
19. Gasparrini A, Guo, Y, Hashizume M *et al.* 2015: Mortality risk attributable to high and low ambient temperature: a multicountry observational study. *The Lancet*. DOI:10.1016/S0140-6736(14)62114-0. See also <https://www.science.dailymail.com/releases/2015/05/150520193831.htm> Accessed June 12, 2018.
20. Coyle, H. 2018: 48,000 Brits dead after worst winter in 42 years. *The Daily Star*. https://www.dailystar.co.uk/news/latest-news/694368/flu-winter-death-cold-fatalities?utm_source=CCNet+Newsletter&utm_campaign=1b394a49be-EMAIL_CAMPAIGN_2018_04_09&utm_medium=email&utm_term=0_fe4b2f45ef-1b394a49be-20143845 April 7, 2018; accessed June 12, 2018.
21. Poulter, S. 2017: Campaigners demand urgent cuts to power bill after number of winter deaths among the elderly rise by 40%. *The Daily Mail* <http://www.dailymail.co.uk/news/article-5109511/Calls-cut-power-bills-winter-deaths-rise-40.html#ixzz4zHs6iwle> November 24, 2017; accessed June 12, 2018.
22. Poulter, S. 2017: Campaigners demand urgent cuts to power bill after number of winter deaths among the elderly rise by 40%. *The Daily Mail* <http://www.dailymail.co.uk/news/article-5109511/Calls-cut-power-bills-winter-deaths-rise-40.html#ixzz4zHs6iwle> November 24, 2017; accessed June 12, 2018.
23. Preston, S. H. 2007: The changing relation between mortality and level of economic development. *International Journal of Epidemiology*. 36 (3): 484–90. PMID 17550952. DOI:10.1093/ije/dym075
24. Roser, M. 2018: Life Expectancy. Published online at OurWorldInData.org. <https://ourworldindata.org/life-expectancy> Accessed June 12, 2018.
25. Lewis, N. and Curry, J. 2018: The impact of recent forcing and ocean heat uptake data on estimates of climate sensitivity. *AMS Journal of Climate*. <https://doi.org/10.1175/JCLI-D-17-0667.1> https://journals.ametsoc.org/doi/10.1175/JCLI-D-17-0667.1?utm_source=CCNet+Newsletter&utm_campaign=a24cafd790-EMAIL_CAMPAIGN_2018_04_24&utm_medium=email&utm_term=0_fe4b2f45ef-a24cafd790-20143845 Accessed June 12, 2018.
26. Tol, R. 2010: The impact of climate change and its policy implications. In Moran and Roskam: *Climate Change: The Facts*. IPA, Melbourne, 68–77.
27. CO2 Coalition 2016: *A Primer on Carbon Dioxide and Climate*. <http://co2coalition.org/publications/a-primer-on-carbon-dioxide-and-climate/> 2016; accessed June 12, 2018.
28. Goklany, I. 2014: Unhealthy Exaggeration. *The Global Warming Policy Foundation* <http://www.thegwpcf.org/indur-goklany-unhealthy-exaggeration/> Accessed June 12, 2018.
29. Easterling, D.R., Horton, B., Jones, P.D., Peterson, T.C., Karl, T.R. *et al.* 1997: Maximum and minimum temperature trends for the globe. *Science* 277: 364–367.



30. Knappenberger, P. C., Michaels, P. E. and Davis, R. E. 2001: The nature of observed climate changes across the United States during the 20th century. *Climate Research* 17: 45-53.
31. Alexander, L. V. *et al.* 2006: Global observed changes in daily climate extremes of temperature and precipitation. *J. Geophys. Res.* 111.
32. Robeson, S.M. 2002: Relationships between mean and standard deviation of air temperature: implications for global warming. *Climate Research* 22: 205–213.
33. Braga, A.L.F., Zanobetti, A., and Schwartz, J. 2002: The effect of weather on respiratory and cardiovascular deaths in 12 U.S. cities. *Environmental Health Perspectives* 110: 859–863.
34. Tam, W.W.S., Wong, T.W., Chair, S.Y., and Wong, A.H.S. 2009: Diurnal temperature range and daily cardiovascular mortalities among the elderly in Hong Kong. *Archives of Environmental and Occupational Health* 64: 202–206.
35. Cao, J., Cheng, Y., Zhao, N., Song, W., Jiang, C., Chen, R., and Kan, H. 2009: Diurnal temperature range is a risk factor for coronary heart disease death. *Journal of Epidemiology* 19: 328–332.
36. Ge, W.Z., Xu, F., Zhao, Z.H., Zhao, J.Z., and Kan, H.D. 2013. Association between diurnal temperature range and respiratory tract infections. *Biomedical and Environmental Sciences* 26: 222–225.
37. Baena-Cagnani, C. and Badellino, H. 2011: Diagnosis of allergy and asthma in childhood. *Current Allergy and Asthma Reports* 11: 71–77.
38. Beasley, R., Crane, J., Lai, C.K., Pearce, N. 2000: Prevalence and etiology of asthma. *Journal of Allergy and Clinical Immunology* 105:466–472.
39. Centers for Disease Control and Prevention, Allergens. <https://www.cdc.gov/climateandhealth/effects/allergen.htm> December 11, 2014; accessed June 12, 2018.
40. Liu, A. H. 2007: Hygiene theory and allergy and asthma prevention. *Paediatric and Neonatal Epidemiology* 21: 3: 2-7.
41. Kozyrskyj, A. L. *et al.* 2007: Increased Risk of Childhood Asthma From Antibiotic Use in Early Life. *Chest Journal* 131 (6): 1753–1759.
42. Ewaschuk, J.B. *et al.* 2011: Effect of pasteurization on immune components of milk: implications for feeding preterm infants. *Applied Physiology, Nutrition and Metabolism* 36(2): 175-182 <https://doi.org/10.1139/h11-008>
43. Loss, G. *et al.* 2011: The protective effect of farm milk consumption on childhood asthma and atopy: The GABRIELA study. *The Journal of Allergy and Clinical Immunology* 128:4; 766-773. DOI: <https://doi.org/10.1016/j.jaci.2011.07.048>
44. Wayne, P., Foster, S., Connolly, J. *et al.* 2002: Production of allergenic pollen by ragweed (*Ambrosia artemisiifolia* L.) is increased in CO₂-enriched atmospheres. *Annals of Allergy, Asthma, and Immunology* 88: 279–282.
45. Lee, Y.S., Dickinson, D.B., Schlager, D., and Velu, J.G. 1979: Antigen E content of pollen from individual plants of short ragweed (*Ambrosia artemisiifolia*). *Journal of Allergy and Clinical Immunology* 63: 336–339.
46. Weber, R.W. 2002: Mother Nature strikes back: global warming, homeostasis, and implications for allergy. *Annals of Allergy, Asthma & Immunology* 88: 251–252.
47. Frei, T. and Gassner, E. 2008: Trends in prevalence of allergic rhinitis and correlation with pollen counts in Switzerland. *International Journal of Biometeorology* 52: 841–847.
48. D'Amato, G., Cecchi, L., Bonini, S., Nunes, C. *et al.* 2007: Allergenic pollen and pollen allergy in Europe. *Allergy* 62: 976–990.
49. Jato, V., Rodriguez-Rajo, F.J., Seijo, M.C., and Aira, M.J. 2009: Poaceae pollen in Galicia (N.W. Spain): characterization and recent trends in atmospheric pollen season. *International Journal of Biometeorology* 53: 333–344.
50. Xu, Z., Huang, C., Su, H., Turner, L.R., Qiao, Z. and Tong, S. 2013: Diurnal temperature range and childhood asthma: a time-series study. *Environmental Health* 12: 10.1186/1476-069X-12-12.
51. Vocks, E., Busch, R., Fröhlich, C., Borelli, S., Mayer, H. and Ring, J. 2001: Influence of weather and climate on subjective symptom intensity in atopic eczema. *International Journal of Biometeorology* 45: 27–33.




52. Byremo, G., Rod, G. and Carlsen, K.H. 2006: Effect of climatic change in children with atopic eczema. *Allergy* 61: 1403–1410.
53. GBD 2015 Mortality and Causes of Death, Collaborators. 2016: Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388 (10053): 1459–1544.
54. Shea, K.M. 2007: Global climate change and children's health. *Pediatrics* 120: 1359–1367.
55. Xu, Z., Huang, C., Turner, L.R., Su, H., Qiao, Z., and Tong, S. 2013: Is diurnal temperature range a risk factor for childhood diarrhea? *PLoS One* 8: e64713.
56. Charles, R.C. and Ryan, E.T. 2011: Cholera in the 21st century. *Current Opinion in Infectious Diseases* 24 (5): 472–7. doi:10.1097/QCO.0b013e32834a88af
57. Vachon, D. 2015: Father of Modern Epidemiology. University of California, Los Angeles, Department of Epidemiology <http://www.ph.ucla.edu/epi/snow/fatherofepidemiology.html> Accessed June 12, 2018.
58. Colwell, R. 1996: Global Climate and Infectious Disease: The Cholera Paradigm. *Science* 274:2025–31.
59. Anderson, Christopher 1991: Cholera Epidemic Traced to Risk Miscalculation. *Nature* 354: 255
60. Hajat, S. and Haines, A. 2002: Associations of cold temperatures with GP consultations for respiratory and cardiovascular disease amongst the elderly in London. *International Journal of Epidemiology* 31: 825–830.
61. Keatinge, W.R. and Donaldson, G.C. 2001: Mortality related to cold and air pollution in London after allowance for effects of associated weather patterns. *Environmental Research* 86: 209–216.
62. Donaldson, G.C. 2006: Climate change and the end of the respiratory syncytial virus season. *Clinical Infectious Diseases* 42: 677–679.
63. Carder, M., McNamee, R., Beverland, I., Elton, R., Cohen, G.R., Boyd, J., and Agius, R.M. 2005: The lagged effect of cold temperature and wind chill on cardiorespiratory mortality in Scotland. *Occupational and Environmental Medicine* 62: 702–710.
64. Nafstad, P., Skrondal, A. and Bjertness, E. 2001: Mortality and temperature in Oslo, Norway. 1990–1995. *European Journal of Epidemiology* 17: 621–627.
65. Gouveia, N., Hajat, S. and Armstrong, B. 2003: Socioeconomic differentials in the temperature-mortality relationship in Sao Paulo, Brazil. *International Journal of Epidemiology* 32: 390–397.
66. Song, G., Chen, G., Jiang, L., Zhang, Y., Zhao, N., Chen, B., and Kan, H. 2008: Diurnal temperature range as a novel risk factor for COPD death. *Respirology* 13: 1066–1069.
67. Hajat, S. and Haines, A. 2002: Associations of cold temperatures with GP consultations for respiratory and cardiovascular disease amongst the elderly in London. *International Journal of Epidemiology* 31: 825–830.
68. Enquesselassie, F., Dobson, A.J., Alexander, H.M. and Steele, P.L. 1993: Seasons, temperature and coronary disease. *International Journal of Epidemiology* 22: 632–636.
69. Nafstad, P., Skrondal, A. and Bjertness, E. 2001: Mortality and temperature in Oslo, Norway, 1990–1995. *European Journal of Epidemiology* 17: 621–627.
70. Kloner, R.A., Poole, W.K. and Perritt, R.L. 1999: When throughout the year is coronary death most likely to occur? A 12-year population-based analysis of more than 220,000 cases. *Circulation* 100: 1630–1634.
71. Kovats, R.S., Hajat, S. and Wilkinson, P. 2004: Contrasting patterns of mortality and hospital admissions during hot weather and heat waves in Greater London, UK. *Occupational and Environmental Medicine* 61: 893–898.
72. Green, M.S., Harari, G. and Kristal-Boneh, E. 1994: Excess winter mortality from ischaemic heart disease and stroke during colder and warmer years in Israel. *European Journal of Public Health* 4: 3–11.
73. Braga, A.L.F., Zanobetti, A. and Schwartz, J. 2002: The effect of weather on respiratory and cardiovascular deaths in 12 U.S. cities. *Environmental Health Perspectives* 110: 859–863.



74. Cagle, A. and Hubbard, R. 2005: Cold-related cardiac mortality in King County, Washington, USA 1980–2001. *Annals of Human Biology* 32: 525–537.
75. Nakaji, S., Parodi, S., Fontana, V. *et al.* 2004: Seasonal changes in mortality rates from main causes of death in Japan (1970–1999). *European Journal of Epidemiology* 19: 905–913.
76. Feigin, V.L., Nikitin, Yu.P. *et al.* 2000: A population-based study of the associations of stroke occurrence with weather parameters in Siberia, Russia (1982–92). *European Journal of Neurology* 7: 171–178.
77. Hong, Y-C., Rha, J-H., Lee, J-T., Ha, E-H., Kwon, H-J., and Kim, H. 2003: Ischemic stroke associated with decrease in temperature. *Epidemiology* 14: 473–478.
78. Chang, C.L. *et al.* 2004: Lower ambient temperature was associated with an increased risk of hospitalization for stroke and acute myocardial infarction in young women. *J. Clin. Epidemiology* 57: 749–757.
79. Gill, R.S., Hambridge, H.L. *et al.* 2012: Falling temperature and colder weather are associated with an increased risk of Aneurysmal Subarachnoid Hemorrhage. *World Neurosurgery* 79: 136–142.
80. Keatinge, W.R. and Donaldson, G.C. 2004: The impact of global warming on health and mortality. *Southern Medical Journal* 97: 1093–1099.
81. Bosello, F., Roson, R. and Tol, R.S.J. 2006: Economy-wide Estimates of the Implications of Climate Change: Human Health. *Ecological Economics* 58(3): 579–91.
82. Gasparrini, A. *et al.* 2017: Projections of temperature-related excess mortality under climate change scenarios. *The Lancet Planetary Health*. DOI: [http://dx.doi.org/10.1016/S2542-5196\(17\)30156-0](http://dx.doi.org/10.1016/S2542-5196(17)30156-0) [https://www.thelancet.com/journals/lanplh/article/PIIS2542-5196\(17\)30156-0/fulltext?elsca1=tlxpr](https://www.thelancet.com/journals/lanplh/article/PIIS2542-5196(17)30156-0/fulltext?elsca1=tlxpr) November 13, 2017; accessed June 12, 2018.
83. Cass, O. 2018: Doomsday Climate Scenarios Are a Joke. *Wall Street Journal* https://www.wsj.com/articles/doomsday-climate-scenarios-are-a-joke-1520800377?utm_source=CCNet+Newsletter&utm_campaign=2aef53c7cc-EMAIL_CAMPAIGN_2018_03_12&utm_medium=email&utm_term=0_fe4b2f45ef-2aef53c7cc-20143845 March 11, 2018; accessed June 12, 2018.
84. Christidis, N., Donaldson, G.C., and Stott, P.A. 2010: Causes for the recent changes in cold- and heat-related mortality in England and Wales. *Climatic Change* 102: 539–553.
85. Urban Analyst, 77 percent of Australia’s population growth over last decade occurred in capital cities: ABS data, <http://www.urbananalyst.com/in-the-news/australia/4506-77-per-cent-of-australia-s-population-growth-over-last-decade-occurred-in-capital-cities-abs-data.html>, July 31, 2017; accessed June 12, 2018.
86. Deschenes, O. and Moretti, E. 2009: Extreme weather events, mortality, and migration. *The Review of Economics and Statistics* 91: 659–681.
87. World Health Organization, Vector-borne Diseases. <http://www.who.int/en/news-room/fact-sheets/detail/vector-borne-diseases> October 31, 2017; accessed June 12, 2018.
88. Intergovernmental Panel on Climate Change (IPCC)-II 2013: Chapter 11, Human Health, Working Group II, IPCC Fifth Assessment Report, March 28, 2013, p. 3.
89. Rogers, D.J. and Randolph, S.E. 2000: The global spread of malaria in a future, warmer world. *Science* 289: 1763–66.
90. Reiter, P. 2000: From Shakespeare to Defoe: malaria in England in the Little Ice Age. *Emerging Infectious Diseases* 6 (1):1-11.
91. Dobson, M. 1989: History of malaria in England. *J. R. Soc. Med.* 82:3-7.
92. Kuhn, K.G., Campbell-Lendrum, D.H., Armstrong, B., and Davies, C.R. 2003: Malaria in Britain: Past, present, and future. *Proceedings of the National Academy of Science, USA* 100: 9997–10001.
93. Hulden, L. and Hulden, L. 2009: The decline of malaria in Finland—the impact of the vector and social variables. *Malaria Journal* 8: 10.1186/1475-2875-8-94.
94. Bruce-Chwatt, L. and de Zulueta, J. 1980: *The rise and fall of malaria in Europe, a historico-epidemiological study*. Oxford University.





95. The History of Malaria 2011: <http://www.malaria.com/overview/malaria-history>; accessed 22 June, 2018.
96. Brierly, W. B. 1944: Malaria and socio-economic conditions in Mississippi. *Social Forces* 23;1: 451-59.
97. CDC 1999: Control of infectious diseases 1900-1999. *JAMA* 282 (11):1029-32. [www.http://jama.ama-assn.org](http://jama.ama-assn.org).
98. Childs, D.Z., Cattadori, I.M. *et al.* 2006: Spatiotemporal patterns of malaria incidence in northern Thailand. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 100: 623-631.
99. World Health Organization 2013: World Malaria Report 2013, Annex 6B. http://www.who.int/malaria/publications/world_malaria_report_2013/en/ Accessed June 12, 2018.
100. Jackson, M.C., Johansen, L., Furlong, C., Colson, A., and Sellers, K.F. 2010: Modelling the effect of climate change on prevalence of malaria in western Africa. *Statistica Neerlandica* 64: 388-400.
101. Hay, S.I., Cox, J., Rogers, D.J., Randolph, S.E., Stern, D.I., Shanks, G.D., Myers, M.F., and Snow, R.W. 2002. Climate change and the resurgence of malaria in the East African highlands. *Nature* 415: 905-909.
102. Shanks, G.D., Biomndo, K., Hay, S.I., and Snow, R.W. 2000: Changing patterns of clinical malaria since 1965 among a tea estate population located in the Kenyan highlands. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 94: 253-255.
103. Small, J., Goetz, S.J., and Hay, S.I. 2003: Climatic suitability for malaria transmission in Africa, 1911-1995. *Proceedings of the National Academy of Sciences USA* 100: 15, 341-15, 345.
104. Rogers, D.J. and Randolph, S.E. 2006: Climate change and vector-borne diseases. *Advances in Parasitology* 62: 345-381.
105. Gething, P. W., Smith, D. L., Patil, A. P., Tatem, A. J., Snow, R. W., Hay, S. I. 2010: Climate change and the global malaria recession. Journal name: *Nature* Volume: 465: Pages: 342-345 Date published: doi:10.1038/nature09098; <https://www.nature.com/articles/nature09098> Accessed June 12, 2018.
106. Haque, U., Hashizume, M., Glass, G.E., *et al.* 2010: The role of climate variability in the spread of malaria in Bangladeshi highlands. *PLoS ONE* 5: 10.1371/journal.pone.0014341.
107. NASA, Carbon Dioxide Fertilization Greening Earth, Study Finds. April 26, 2016 <https://www.nasa.gov/feature/goddard/2016/carbon-dioxide-fertilization-greening-earth>; accessed June 12, 2018.
108. Transer, F. C., Sharp, B. and Sueur, D. 2003: Potential Effect of Climate Change on Malaria Transmission in Africa. *Royal Society of Tropical Medicine & Hygiene* 97: 129-32.
109. Reiter P. 1998: Global-warming and vector-borne disease in temperate regions and at high altitude. *Lancet* 1998; 351: 839-40.
110. Béguin, A., Hales, S., Rocklöv, J., Åström, C. *et al.* 2011: The opposing effects of climate change and socio-economic development on the global distribution of malaria. *Global Environmental Change* 21: 1209-1214.
111. Goklany, I. 2004: Climate Change and Malaria. *Science* 306: 57, October 1, 2014.
112. Robinson, M.C. 1955: An epidemic of virus disease in Southern Province, Tanganyika Territory, in 1952-53. I. Clinical features. *Trans. R. Soc. Trop. Med. Hyg.* 49: 28-32.
113. Reiter, P. 2009: Yellow fever and dengue: a threat for Europe. *Euro. Surveill.* 15: pii=19509
114. Almeida, A.P.G., Gonçalves, Y.M., Novo, M.T., Sousa, C.A., Melim, M., Gracio, A.J. 2007: Vector monitoring of *Aedes aegypti* in the Autonomous Region of Madeira, Portugal. *Euro. Surveill.* 12: pii=3311.
115. Iunicheva, I.V., Riabova, T.E., Bezzhonova, O.V., Ganushkina, L.A., *et al.* 2008: First evidence for breeding *Aedes aegypti* L in the area of Greater Sochi and in some towns of Abkhasia. *Med. Parazitol.* 3: 40-43.
116. Wikipedia: Global map of the predicted distribution of *Aedes aegypti* in 2015. [https://en.wikipedia.org/wiki/Aedes_aegypti#/media/File:Global_Aedes_aegypti_distribution_\(e08347\).png](https://en.wikipedia.org/wiki/Aedes_aegypti#/media/File:Global_Aedes_aegypti_distribution_(e08347).png) June 30, 2015; accessed June 12, 2018.
117. Wikipedia: Global map of the predicted distribution of *Aedes aegypti* in 2015. [https://en.wikipedia.org/wiki/Aedes_aegypti#/media/File:Global_Aedes_aegypti_distribution_\(e08347\).png](https://en.wikipedia.org/wiki/Aedes_aegypti#/media/File:Global_Aedes_aegypti_distribution_(e08347).png) June 30, 2015; accessed June 12, 2018.

- 
118. Burt, F.J., Rolpf, M.S., Rulli, N.E, *et al.* 2012: Chikungunya: a re-emerging virus. *Lancet*. 379: 662-671.
 119. Scholte, E-J. and Schaffner, F: 2007: *Waiting for the tiger: establishment and spread of the Aedes albopictus mosquito in Europe. Emerging pests and vector-borne diseases in Europe*. Edited by: Takken, W., Knols, B.G.J. Wageningen: Wageningen Academic Publishers, 241-60.
 120. Hii, Y.L., Rocklöv, J., Ng N, Tang, C.S, Pang, F.Y., Sauerborn, R. 2009: Climate variability and increase in intensity and magnitude of dengue incidence in Singapore. *Glob. Health Action* 11: 2 doi:10.3402/gha.v2i0.2036
 121. Hanson, S. M. and Craig, G. B. 1995: *Aedes albopictus* (Diptera: Culcidae) Eggs: Field Survivorship During Northern Indiana Winters. *Journal of Medical Entomology* 32 (5): 599–604. DOI:10.1093/jmedent/32.5.599
 122. Romi, R., Severini, F., Toma., L. 2006: Cold acclimation and overwintering of female *Aedes albopictus* in Roma. *Journal of the American Mosquito Control Association* 22 (1): 149–51. DOI:10.2987/8756-971X(2006)22[149:CAAOOF]2.0.CO;2. PMID 16646341
 123. Adhami. J. and Reiter, P. 1998: Introduction and establishment of *Aedes* (Stegomyia) *albopictus* Skuse (Diptera: Culicidae) in Albania. *J. Am. Mosq. Control Assoc.* 14: 340-343.
 124. Sabatini, A., Raineri, V., Trovato, V.G., Coluzzi, M. 1990: *Aedes albopictus* in Italia e possibile diffusione della specie nell' area Mediterranea. *Parassitologia* 32: 301-304.
 125. Rezza, G., Nicoletti, L., Angelini, R., Romi, R., Finarelli, A.C., Panning, M. 2007: Infection with Chikungunya virus in Italy: an outbreak in a temperate region. *Lancet* 370: 1840-1846.
 126. Staples, J.E., Fischer, M. 2014: Chikungunya Virus in the Americas — What a Vectorborne Pathogen Can Do. *N. Engl. J. Med.* 371 (10): 887–9. DOI:10.1056/NEJMp1407698 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4624217/> September 4, 2014; accessed June 12, 2018.
 127. Centers for Disease Control and Prevention, Chikungunya virus in the United States. January 25, 2018: <https://www.cdc.gov/chikungunya/geo/united-states.html> January 25 accessed June 12, 2018.
 128. Wikimedia, Albopictusdistribution2007: https://upload.wikimedia.org/wikipedia/commons/0/02/Albopictus_distribution_2007.png; accessed June 12, 2018.
 129. Fischer, D., Thomas, S.M., Suk, J.E., Sudre, B., Hess, A. Tjaden, N.B., Beierkuhnlein, N. and Semenza, J.C. 2013: Climate change effects on Chikungunya transmission in Europe: geospatial analysis of vector's climatic suitability and virus' temperature requirements. *International Journal of Health Geographics* 12:51 <https://doi.org/10.1186/1476-072X-12-51>; <https://ij-healthgeographics.biomedcentral.com/articles/10.1186/1476-072X-12-51> November 12, 2013; accessed June 12, 2018.
 130. Tuchman, N.C., Wahtera, K.A., Wetzel, R.G., Russo, N.M. *et al.* 2003: Nutritional quality of leaf detritus altered by elevated atmospheric CO₂: effects on development of mosquito larvae. *Freshwater Biology* 48: 1432–1439.
 131. Kyle, J.L. and Harris, E. 2008: Global spread and persistence of dengue. *Annual Review of Microbiology* 62: 71–92.
 132. Wilder-Smith, A. and Gubler, D.J. 2008: Geographic expansion of Dengue: The impact of international travel. *Medical Clinics of North America* 92: 1377–1390.
 133. Beatty, M.E., Letson, G.W. and Margolis, H.S. 2008: Estimating the global burden of dengue. Proceedings of the 2nd International Conference on Dengue and Dengue Haemorrhagic Fever. Phuket, Thailand.
 134. Naish, S., Dale, P., Mackenzie, J.S. *et al.* 2014: Climate change and dengue: a critical and systematic review of quantitative modelling approaches. *BMC Infectious Diseases* 14:167 <https://doi.org/10.1186/1471-2334-14-167> <https://bmcinfectdis.biomedcentral.com/articles/10.1186/1471-2334-14-167> March 26, 2014; accessed June 12, 2018.
 135. Climate Change Guide, Dengue Fever and Climate Change. <http://www.climate-change-guide.com/dengue-fever.html> ND; accessed June 12, 2018.
 136. Kyle, J.L. and Harris, E. 2008: Global spread and persistence of dengue. *Annual Review of Microbiology* 62: 71–92.
 137. Nabi, S.A. and Qader, S.S. 2009: Is global warming likely to cause an increased incidence of malaria? *Libyan Journal of Medicine* 4: 18–22.




138. Centers for Disease Control and Prevention, Dengue and Climate: September 27, 2012 <https://www.cdc.gov/dengue/entomologyecology/climate.html#travelers> accessed June 12, 2018.
139. Russell, R.C., Currie, B.J., Lindsay, M.D., *et al.* 2009: Dengue and climate change in Australia: predictions for the future should incorporate knowledge from the past. *Medical Journal of Australia* 190: 265–268.
140. Russell, R.C. 2009: Mosquito-borne disease and climate change in Australia: time for a reality check. *Australian Journal of Entomology* 48: 1–7.
141. Wilder-Smith, A. and Gubler, D.J. 2008: Geographic expansion of Dengue: The impact of international travel. *Medical Clinics of North America* 92: 1377–1390.
142. Moore, T. G. 1998: *Climate of Fear*, Cato Institute, Washington DC p. 78.
143. Reiter, P., Lathrop, S., Bunning, M., *et al.* 2003: Texas lifestyle limits transmission of Dengue virus. *Emerging Infectious Diseases* 9: 86–89.
144. Centers for Disease Control and Prevention, Dengue and Climate. <https://www.cdc.gov/dengue/entomologyecology/climate.html> September 27, 2012; accessed June 12, 2018.
145. Takeda, Takeda's Dengue Vaccine Candidate Associated with Reduced Incidence of Dengue in Children and Adolescents; New 18-Month Interim Phase 2 Data Published in *The Lancet Infectious Diseases*. <https://www.takeda.com/newsroom/newsreleases/2017/takedas-dengue-vaccine/> November 6, 2017; accessed June 12, 2018.
146. Wikipedia: *Aedes aegypti*. https://en.wikipedia.org/wiki/Aedes_aegypti April 4, 2018; accessed June 12, 2018.
147. Wikipedia: Sterile insect technique. https://en.wikipedia.org/wiki/Sterile_insect_technique June 2, 2018; accessed June 12, 2018.
148. Wikipedia: *Wolbachia*. <https://en.wikipedia.org/wiki/Wolbachia> May 15, 2018; accessed June 12, 2018.
149. Epstein, P.R. 2001: West Nile virus and the climate. National Center for Biotechnology Information, National Library of Medicine, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3456354/> June 2001; accessed June 12, 2018.
150. Walsh, B. 2014: Thanks to Climate Change, West Nile Virus Could Be Your New Neighbor. *Time* <http://time.com/11683/west-nile-virus-climate-change/> Feb. 28, 2014; accessed June 12, 2018.
151. Centers for Disease Control and Prevention, West Nile virus disease cases and deaths reported to CDC by year and clinical presentation, 1999–2016. <https://www.cdc.gov/westnile/statsmaps/cumMapsData.html#three> July 15, 2016; accessed June 12, 2018.
152. Wikipedia: West Nile virus in the United States. https://en.wikipedia.org/wiki/West_Nile_virus_in_the_United_States December 18, 2018; accessed June 12, 2018.
153. McNeill, J. R. 2004: Yellow Jack and Geopolitics: Environment, Epidemics, and the Struggles for Empire in the American Tropics, 1650–1825. *OAH Magazine of History*. 18 (3): 9–13. DOI:10.1093/maghis/18.3.9 <https://academic.oup.com/maghis/article-abstract/18/3/9/1034950>
154. Wikipedia: 1793 Philadelphia yellow fever epidemic. https://en.wikipedia.org/wiki/1793_Philadelphia_yellow_fever_epidemic May 20, 2018; accessed June 12, 2018.
155. Campbell, B. C. ed. 2008: American Disasters: 201 Calamities That Shook the Nation. pp 49–50.
156. Marr, J.S. and Cathey, J.T. 2013: The 1802 Saint-Domingue yellow fever epidemic and the Louisiana Purchase. *Journal of Public Health Management and Practice* 19#1
157. Sawchuck, L.A. and Burke, S. D. A. 1998: Gibraltar's 1804 Yellow Fever Scourge: The Search for Scapegoats. <https://academic.oup.com/jhmas/article-abstract/53/1/3/742694> January 1, 1998; accessed June 12, 2018.
158. Canela Soler, J., Pallarés Fusté, M.R. *et al.* 2008: A mortality study of the last outbreak of yellow fever in Barcelona City (Spain) in 1870. *Gaceta sanitaria / S.E.S.P.A.S.* 23 (4): 295–9. DOI:10.1016/j.gaceta.2008.09.008
159. Coleman, W. 1983: Epidemiological method in the 1860s: yellow fever at Saint-Nazaire. *Bulletin of the history of medicine*. 58 (2): 145–63. PMID 6375767 <https://www.ncbi.nlm.nih.gov/pubmed/6375767> accessed June 12, 2018.

- 
160. Meers, P.D. 1986: Yellow fever in Swansea, 1865. *The Journal of hygiene* 97 (1): 185–91. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2082871/> August 1986; accessed June 12, 2018.
 161. Barrett, A.D., Teuwen, D.E. 2009: Yellow fever vaccine – how does it work and why do rare cases of serious adverse events take place? *Current Opinion in Immunology* 21 (3): 308–13. DOI:10.1016/j.coi.2009.05.018
 162. Lane, D. *et al.* 2017: How Zika upped the ante. *A*Star Research* 8,12-15. www.research.a-star.edu.sg/feature-and-innovation/7740/how-zika-upped-the-ante August 10, 2017; accessed June 12, 2018.
 163. Randolph, S.E. and Rogers, D.J. 2000: Fragile transmission cycles of tick-borne encephalitis virus may be disrupted by predicted climate change. *Proceedings of the Royal Society of London Series B* 267: 1741–1744.
 164. Randolph, S.E. 2001: The shifting landscape of tick-borne zoonoses: tick-borne encephalitis and Lyme borreliosis in Europe. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 356(1411):1045–1056.
 165. Lindgren, E. and Jaensen, T. 2006: Lyme borreliosis in Europe: influences of climate and climate change, epidemiology, ecology and adaptation measures. World Health Organization http://www.euro.who.int/_data/assets/pdf_file/0006/96819/E89522.pdf 2006; accessed June 12, 2018.
 166. Sumilo, D., Bormane, A., Asokliene, L., *et al.* 2008: Socio-economic factors in the differential upsurge of tick-borne encephalitis in Central and Eastern Europe. *Reviews in Medical Virology* 18: 81–95.
 167. Randolph, S.E. 2010: To what extent has climate change contributed to the recent epidemiology of tick-borne diseases? *Veterinary Parasitology* 167: 92–94.
 168. Williams, Carolyn. 2007: *Infectious Disease Epidemiology: Theory and Practice* (2nd ed.). Sudbury, Mass.: Jones and Bartlett Publishers. p. 447. ISBN 9780763728793; Wikipedia: https://en.wikipedia.org/wiki/Old_Lyme,_Connecticut
 169. Shapiro, E.D. 2014: Clinical Practice. Lyme Disease. (PDF). *N. Engl. J. Med.* 370 (18): 1724–31. https://web.archive.org/web/20161019142422/http://portal.mah.harvard.edu/templatesnew/departments/MTA/Lyme/uploaded_documents/NEJMcp1314325.pdf May 5, 2014; accessed June 12, 2018.
 170. Lindgren, E. and Jaensen, T. 2006: Lyme borreliosis in Europe: influences of climate and climate change, epidemiology, ecology and adaptation measures. World Health Organization http://www.euro.who.int/_data/assets/pdf_file/0006/96819/E89522.pdf 2006; accessed June 12, 2018.
 171. Pritt, B.S., Mead, P.S., Johnson, D.K., Neitzel, D.F. *et al.* 2016: Identification of a novel pathogenic *Borrelia* species causing Lyme borreliosis with unusually high spirochaetaemia: a descriptive study. *Lancet Infect. Dis.* 16: 556–564. doi:10.1016/S1473-3099(15)00464-8; [https://www.thelancet.com/journals/laninf/article/PIIS1473-3099\(15\)00464-8/fulltext](https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(15)00464-8/fulltext) February 5, 2016; accessed June 12, 2018; Wikipedia: *Borrelia mayonii* https://en.wikipedia.org/wiki/Borrelia_mayonii May 3, 2018; accessed June 12, 2018.
 172. Clark, K. 2004: *Borrelia* Species in Host-Seeking Ticks and Small Mammals in Northern Florida. *J. Clin. Microbiol.* 42 (11): 5076–86. DOI:10.1128/JCM.42.11.5076-5086.2004 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC525154/> November 2004; accessed June 12, 2018.
 173. Süss, J., Klaus, C., Gerstengarbe, F.W., Werner, P.C. 2008: What Makes Ticks Tick? Climate Change, Ticks, and Tick-Borne Diseases. *J Travel Med.* 15 (1): 39–45. DOI:10.1111/j.1708-8305.2007.00176 <https://academic.oup.com/jtm/article/15/1/39/1849257> January 21, 2008; accessed June 12, 2018.
 174. Centers for Disease Control and Prevention, *Lyme Disease: What You Need to Know*. <https://www.cdc.gov/lyme/resources/brochure/lymediseasebrochure.pdf> ND; accessed June 12, 2018.
 175. Subak, S. 2003: Effects of Climate on Variability in Lyme Disease Incidence in the Northeastern United States. *American Journal of Epidemiology* 105 (6): 531–538.
 176. Jones, C.J. and Kitron, U.D. 2000: Populations of *Ixodes scapularis* (Acari: Ixodidae) Are Modulated by Drought at a Lyme Disease Focus in Illinois. *Journal of Medical Entomology*, 37 (3): 408–415.
 177. Centers for Disease Control and Prevention, *Lyme Disease: What You Need to Know*. <https://www.cdc.gov/lyme/resources/brochure/lymediseasebrochure.pdf> ND; accessed June 12, 2018.




178. Brownstein, J.S., Holford, T.R. and Fish, D. 2005: Effect of Climate Change on Lyme Disease Risk in North America. *Ecohealth* 2(1): 38–46. DOI:10.1007/s10393-004-0139-x; <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2582486/>
179. Lindgren, E. and Jaensen, T. 2006: Lyme borreliosis in Europe: influences of climate and climate change, epidemiology, ecology and adaptation measures. World Health Organization http://www.euro.who.int/__data/assets/pdf_file/0006/96819/E89522.pdf 2006; accessed June 12, 2018.
180. Zeman, P. and Januska, J. 1999: Epizootic background of dissimilar distribution of human cases of Lyme borreliosis and tick-borne encephalitis in a joint endemic area. *Comparative Immunology, Microbiology and Infectious Disease*, 22: 247–260.
181. Daniel, M, *et al.* 2004: An attempt to elucidate the increased incidence of tick-borne encephalitis and its spread to higher altitudes in the Czech Republic. *International Journal of Medical Microbiology* 293(37):55–62.
182. Lindgren, E. and Gustafson, R. 2001: Tick-borne encephalitis in Sweden and climate change. *Lancet* 358:16–18.
183. Lindgren, E. and Jaensen, T. 2006: Lyme borreliosis in Europe: influences of climate and climate change, epidemiology, ecology and adaptation measures. World Health Organization http://www.euro.who.int/__data/assets/pdf_file/0006/96819/E89522.pdf 2006; accessed June 12, 2018.
184. Lindgren, E. and Jaensen, T. 2006: Lyme borreliosis in Europe: influences of climate and climate change, epidemiology, ecology and adaptation measures. World Health Organization http://www.euro.who.int/__data/assets/pdf_file/0006/96819/E89522.pdf 2006; accessed June 12, 2018.
185. World Health Organization: Leishmaniasis. <http://www.who.int/en/news-room/fact-sheets/detail/leishmaniasis> March 14, 2018; accessed June 12, 2018.
186. Desjeux, P. 2001: Worldwide increasing risk factors for leishmaniasis. *Med. Microbiol. Immunol.* (Berlin), 190, 77-79.
187. Postigo, J.A. 2010: Leishmaniasis in the World Health Organization Eastern Mediterranean Region. *Int J Antimicrob. Agents* 36 (Suppl 1)
188. Pratlong, F., Dereure, J., Bucheton, B., El-Saf, S., Dessein, A., Lanotte, G. *et al.* 2001: Sudan: the possible original focus of visceral leishmaniasis. *Parasitology* 122: 599-605.
189. Gonzalez, C., Wang, O., Strutz, S.E., *et al.* 2010: Climate change and risk of leishmaniasis in North America: predictions from ecological niche models of vector and reservoir species. *P LoS Negl. Trop. Dis.* 4 (1): e585.
190. González, Camila *et al.* 2010: Climate Change and Risk of Leishmaniasis in North America: Predictions from Ecological Niche Models of Vector and Reservoir Species. *PLOS* <https://doi.org/10.1371/journal.pntd.0000585> Figure 1: <http://journals.plos.org/plosntds/article/figure?id=10.1371/journal.pntd.0000585.g001>; accessed June 12, 2018.
191. Weather.com, The Storm that Killed 300,000. <https://weather.com/storms/hurricane/news/deadliest-cyclone-history-bangladesh-20130605#/1> April 25, 2014; accessed June 12, 2018.
192. Wikipedia: Great Hurricane of 1780. https://en.wikipedia.org/wiki/Great_Hurricane_of_1780 May 3, 2018; accessed June 12, 2018.
193. Wikipedia: 1815 New England hurricane. https://en.wikipedia.org/wiki/1815_New_England_hurricane April 19, 2018; accessed June 12, 2018.
194. Wikipedia: 1821 Norfolk and Long Island hurricane. https://en.wikipedia.org/wiki/1821_Norfolk_and_Long_Island_hurricane April 2, 2018; accessed June 12, 2018.
195. Wikipedia: 1893 New York hurricane. https://en.wikipedia.org/wiki/1893_New_York_hurricane April 22, 2018; accessed June 12, 2018.
196. Nyberg, J., Winter, A., Malmgren, B.A. 2005: *Reconstruction of Major Hurricane Activity*. *Eos Trans. AGU.* 86 (52, Fall Meet. Suppl.): Abstract. P.21C–1597.
197. Nott, J., Haig, J., Neil, H. and Gillieson, D. 2007: Greater frequency variability of landfalling tropical cyclones at centennial compared to seasonal and decadal scales. *Earth and Planetary Science Letters* 255; 367-72. <https://doi.org/10.1016/j.epsl.2006.12.023>

- 
198. Bureau of Meteorology, Government of Australia, Tropical Cyclone Trends. 2018: <http://www.bom.gov.au/cyclone/climatology/trends.shtml>; accessed June 12, 2018.
 199. Haig, J.E-A. and Nott, J. 2016: Solar forcing over the last 1500 years and Australian tropical cyclone activity. *Geophysical Research Letters* 43(6): 2843-2850; <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL068012> March 6, 2016; accessed June 12, 2018.
 200. Australian Bureau of Statistics, 1301.0 - Yearbook Complete, 2008. Feature Article 3: Understanding Natural Hazard Impacts on Australia. <http://www.abs.gov.au/ausstats/abs@.nsf/Previousproducts/1301.0Feature%20Article42008> February 7, 2008; accessed June 12, 2018.
 201. ABC News, Tropical Cyclone Mahina: Bid to have deadly March 1899 weather event upgraded in record books. <http://www.abc.net.au/news/2014-12-26/cyclone-mahina/5964342> December 26, 2014; accessed June 12, 2018.
 202. Euronews, No loss of life as Cyclone Yasi hits Australia. <https://www.youtube.com/watch?v=GjYA-X6-gJc> February 3, 2011; accessed June 12, 2018.
 203. WMO Tropical Meteorology Research Program, Summary Statement on Tropical Cyclones and Climate Change https://web.archive.org/web/20090325193707/http://www.wmo.int/pages/prog/arep/press_releases/2006/pdf/iwtc_summary.pdf 2006; accessed June 12, 2018.
 204. Pielke, R.A., Jr. et al. 2008: *Normalized Hurricane Damage in the United States: 1900–2005* (PDF). *Natural Hazards Review* 9 (1): 29–42. DOI:10.1061/(ASCE)1527-6988(2008)9:1(29) November 6, 2018; accessed June 12, 2018.
 205. Wikipedia: 1926 Miami hurricane https://en.wikipedia.org/wiki/1926_Miami_hurricane February 27, 2018; accessed June 12, 2018.
 206. Wikipedia: Typhoon <https://en.wikipedia.org/wiki/Typhoon> May 28, 2018; accessed June 12, 2018.
 207. Anderson-Berry, L.J. and Weyman, J.C. 2008: Fifth International Workshop on Tropical Cyclones: Topic 5.1: Societal Impacts of Tropical Cyclones. *World Meteorological Organization. National Oceanic and Atmospheric Administration*. December 2002; accessed June 12, 2018.
 208. Dunnavan, G.M. and Diercks, J.W. 1980: An Analysis of Super Typhoon Tip (October 1979). *Monthly Weather Review*. American Meteorological Society. 108 (11): 1915–1923. Bibcode:1980MWRv.108.1915D. DOI:10.1175/1520-0493(1980)108<1915:AAOSTT>2.0.CO;2. ISSN 1520-0493, <https://journals.ametsoc.org/doi/pdf/10.1175/1520-0493%281980%29108%3C1915%3AAAOSTT%3E2.0.CO%3B2> July 29, 1980; accessed June 12, 2018.
 209. Samenow, J. and McNoldy, B. 2013: Super typhoon Haiyan strikes Philippines, among strongest storms ever. *Washington Post*, https://www.washingtonpost.com/news/capital-weather-gang/wp/2013/11/07/super-typhoon-haiyan-closes-in-on-philippines-among-strongest-storms-ever/?noredirect=on&utm_term=.03eeb22dda79 November 7, 2013; accessed June 12, 2018.
 210. Derived from table of typhoons in Wikipedia: Typhoon <https://en.wikipedia.org/wiki/Typhoon> May 28, 2018; accessed June 12, 2018.
 211. <https://www.independent.co.uk/environment/world-deadliest-storms-tornadoes-cyclones-hail-lightning-bangladesh-india-egypt-zimbabwe-a7741261.html>
 212. Johnston, I. 2017: World's deadliest storms from tornadoes to cyclones, lightning and hail. *Independent*. <https://www.independent.co.uk/environment/world-deadliest-storms-tornadoes-cyclones-hail-lightning-bangladesh-india-egypt-zimbabwe-a7741261.html> May 18, 2017; accessed June 12, 2018.
 213. The Conversation, 2017: To understand how storms batter Australia, we need a fresh deluge of data. <https://theconversation.com/to-understand-how-storms-batter-australia-we-need-a-fresh-deluge-of-data-68487> accessed June 12, 2018.
 214. Hermida, L., Sánchez, J.L., López, L., Berthet, C., Dessens, J., Garcia-Ortega, E., and Merino, A. 2013: Climatic Trends in Hail Precipitation in France: Spatial, Altitudinal, and Temporal Variability. *The Scientific World Journal*, Volume 2013, Article ID 494971, 10 pages <http://dx.doi.org/10.1155/2013/494971>, <https://www.hindawi.com/journals/tswj/2013/494971/> 2013; accessed June 12, 2018.




215. Changnon, S.A. and Changnon, D. 2000: *Long-Term Fluctuations in Hail Incidences in the United States*. *J. Climate*, 13: 658–664, [https://doi.org/10.1175/1520-0442\(2000\)013<0658:LTFIHI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<0658:LTFIHI>2.0.CO;2), <https://journals.ametsoc.org/doi/abs/10.1175/1520-0442%282000%29013%3C0658%3ALTFIHI%3E2.0.CO%3B2> February 1, 2000; accessed June 12, 2018.
216. NOAA's National Weather Service Storm Prediction Center, Inflation Adjusted Annual Tornado Running Total. <http://www.spc.noaa.gov/wcm/adj.html> January 5, 2018; accessed June 12, 2018.
217. Lepore, C., Tippet, M.K. and Allen, J.T. 2017: ENSO-based probabilistic forecasts of March–May U.S. tornado and hail activity. *Geophysical Research Letters*, 44;17, 9093–9101 <https://doi.org/10.1002/2017GL074781>
218. NOAA's National Weather Service Storm Prediction Center, Inflation Adjusted Annual Tornado Running Total. <http://www.spc.noaa.gov/wcm/adj.html> January 5, 2018; accessed June 12, 2018.
219. NOAA National Weather Service U.S. Annual Count of Strong to Violent Tornadoes (F3+), 1954 through 2014. <https://www1.ncdc.noaa.gov/pub/data/cmb/images/tornado/clim/EF3-EF5.png> ND; accessed June 12, 2018.
220. White, M. 2012: *The Great Big Book of Horrible Things: The Definitive Chronicle of History's 100 Worst Atrocities*. W. Norton, p. 47. ISBN 9780393081923
221. Demaree, G. R. 2006: The catastrophic floods of February 1784 in and around Belgium – A Little Ice Age event of frost, snow, river ice and floods. *Hydrological Sciences Journal*, 51(5); 878–98.
222. Mudelsee, M. et al. 2006: Trends in flood risk of the river Wierra (Germany) over the last 500 years. *Hydrological Sciences Journal*, 51(5): 818–33.
223. Mitchell, J. K. 2003: European river floods in a changing world. *Risk Analysis*, 23(3):567–74.
224. Yiou, R. et al. 2006: Statistical analysis of floods in Bohemia (Czech Republic) since 1825. *Hydrological Sciences Journal*, 51(5): 930–45.
225. Australian Geographic, Top 10 worst floods in Australia: <http://www.australiangeographic.com.au/topics/history-culture/2012/03/floods-10-of-the-deadliest-in-australian-history> March 8, 2012; accessed June 12, 2018.
226. Bureau of Meteorology, Government of Australia, Known Floods in the Brisbane & Bremer River Basin http://www.bom.gov.au/qld/flood/fld_history/brisbane_history.shtml November 2017; accessed June 12, 2018.
227. Kundzewicz, Z. W. et al. 2005: Trend detection in river flow series: 1. Annual maximum flow. *Hydrological Sciences Journal*, 50(5); 797–810.
228. Svensson, C., Kundzewicz, Z. W. and Maurer, T. 2005: Trend detection in river flow series: 1. Annual maximum flow. *Hydrological Sciences Journal*, 50(5): 811–24. DOI:10.1623/hysj.2005.50.5.811
229. Roger Pielke Jr.'s Blog, Coverage of Extreme Events in the IPCC AR5: http://rogerpielkejr.blogspot.com/2013/10/coverage-of-extreme-events-in-ipcc-ar5.html?utm_source=CCNet+Newsletter&utm_campaign=e4674d7290-EMAIL_CAMPAIGN_2018_04_10&utm_medium=email&utm_term=0_fe4b2f45ef-e4674d7290-20143845 October 3, 2015; accessed June 12, 2018.
230. International Federation of Red Cross and Red Crescent Societies 2015. *World disasters report*. Geneva, Switzerland.
231. Chowdhury, A. M. R., Bhuyia, A. U., Choudhury, A. Y. and Sen, R. 1993: The Bangladesh cyclone of 1991: why so many people died. *Disasters*, 17: 291–304.
232. Haynes, K., Coates, L., de Oliveira, F.D., et al. 2016: An analysis of human fatalities from floods in Australia 1900–2015. *Risk Frontiers*, Macquarie University. https://www.riskfrontiers.com/pdf/flood_fatality_report_final.pdf May 2016; accessed June 12, 2018.
233. Ashley, S. T. and Ashley, W. S. 2008: Flood fatalities in the United States. *Journal of Applied Meteorology and Climatology*, 47: 805–818. <http://commons.lib.niu.edu/bitstream/handle/10843/13369/Ashley,%20S.%20-%20Flood%20Fatalities%20in%20United%20States.pdf;sequence=1> July 5, 2007; accessed June 12, 2018.

- 
234. Haynes, K., Coates, L., de Oliveira, F.D., Gissing, A., Bird, D., van den Honert, R., Radford, D., D'Arcy, R. and Smith, C. 2016: An analysis of human fatalities from floods in Australia 1900-2015. *Risk Frontiers*, Macquarie University. https://www.riskfrontiers.com/pdf/flood_fatalities_report_final.pdf May 2016; accessed June 12, 2018.
 235. Bureau of Meteorology, Government of Australia, Annual Climate Statement 2017: <http://www.bom.gov.au/climate/current/annual/aus/#tabs=Rainfall> March 27, 2018; accessed June 12, 2018.
 236. Bharti, A.R., Nally, J.E., Ricaldi, J.N. *et al.* 2003: Leptospirosis: a zoonotic disease of global importance. *Lancet Infect Dis* 3(12) 757-771.
 237. Lomborg, B. 2007: *Cool It*. Knopf, Borozoi Books, USA p. 83.
 238. To retrieve relevant data, go to the National Climatic Data Center website: <https://www7.ncdc.noaa.gov/CDO/CDODivisionalSelect.jsp>
 239. Meko, D. *et al.* 2007: Medieval drought in the upper Colorado River basin, *Geophysical Research Letters* 34 L10705, doi:10.1029/2007GL029988
 240. Smith T. 2012: Climate change, desertification and migration: Connecting the dots, *Climate Change News*, <http://www.climatechangenews.com/2012/04/27/climate-change-desertification-and-migration-connecting-the-dots/> April 27, 2012; accessed June 12, 2018.
 241. Giannini, A., Saravanin, R. and Chang, P. 2003: Oceanic forcing of Sahel rainfall on interannual to Interdecadal time scales. *Science* 302; 1027-1030. https://iri.columbia.edu/~alesall/pubslst/sciencemag7Nov_gianninietal.pdf
 242. NOAA 2003: <https://www.gfdl.noaa.gov/sahel-drought/> October 9, 2003; accessed June 12, 2018.
 243. Lau, W.K.M., Wu, H.T. and Kim, K.M. 2013: A canonical response of precipitation characteristics to global warming from CMIP5 models, *Geophys. Res. Lett.* 40: 3163–3169. DOI:10.1002/grl.50420.
 244. Sun, F., Roderick, M.L. and Farquhar, G.D. 2012: Changes in the variability of global land precipitation, *Geophys. Res. Lett* 39: L19402. DOI:10.1029/2012GL053369.
 245. Greve, P., Orlowsky, B., Mueller, B. *et al.* 2014: Global assessment of trends in wetting and drying over land, *Nature Geoscience*. DOI:10.1038/ngeo2247.
 246. Royer, D. L. 2001: Stomatal density and stomatal index as indicators of paleoatmospheric carbon dioxide concentration. *Review of Palaeobotany and Palynology*, 114 (1): 1–28.
 247. Ainsworth, E. A. and Long, S P. 2005: What have we learned from fifteen years of free-air CO₂ enrichment FACE? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytologist* 165: 351-372.
 248. Donohue, R.J., Roderick, M.L., McVicar, T.R. and Farquha, G.D. 2013: Carbon dioxide fertilisation has increased maximum foliage cover across the globe's warm, arid environments. *Geophysical Research Letters*. DOI: 10.1002/grl.50563.
 249. Konzmann, M., Gerten, D. and Heinke, J. 2013: Climate impacts on global irrigation requirements under 19 GCMs, simulated with a vegetation and hydrology model. *Hydrological Sciences Journal* 58 (1): 88–105.
 250. Wiltshire, A. *et al.* 2013: The importance of population, climate change and carbon dioxide plant physiological forcing in determining future global water stress. *Global Environmental Change*. 23 (5): 1083–1097.
 251. Schar, C. *et al.* 2004: The role of increasing temperature variability in European summer heatwaves, *Nature* doi: 10.1038/nature02300 <http://www.met.reading.ac.uk/~vidale/papers/SchaerEtAl2004.pdf> accessed June 12, 2018.
 252. Founda, D. and Giannakopoulos, C. 2009: The exceptionally hot summer of 2007 in Athens, Greece—a typical summer in the future climate? *Glob. Planet. Change* 67: 227-36.
 253. Barriopedro, D., Fischer, E., Luterbacher, J., Trigo, R. M. and Garcia-Herrera, R. 2011: The hot summer of 2010: redrawing the temperature record map of Europe. *Science* 332: 220-224.
 254. Coumou, D. and Rahmstorf, S. 2012: A decade of weather extremes. *Nature Climate Change*. 2: 491-496 doi:10.1038/nclimate1452



255. Karoly, D. J. 2009: The recent bushfires and extreme heat wave in southeast Australia. *Bull. Aust. Meteorol. Oceanogr. Soc.* 22: 10–13.
256. Lewis, S. C. and Karoly, D. J. 2013: Anthropogenic contributions to Australia's record summer temperatures of 2013. *Geophys. Res. Lett.*, doi:10.1002/grl.50673.
257. Coumou, D. Robinson, A. and Rahmstorf, S. 2013: Global increase in record-breaking monthly-mean temperatures *Climate Change*, 118: 771–782.
258. Chase, T. N., Wolter, K., Pielke, R. A. Sr. and Rasool, I. 2006: Was the 2003 European summer heat wave unusual in a global context? *Geophysical Research Letters*. 33(5) doi:10.1029/2006GL027470 Was the 2003 European summer heat wave unusual in a global context?
259. Barrier Miner 1896: Thunderstorms and High Temperatures. <https://trove.nla.gov.au/newspaper/article/44159099?zoomLevel=5> January 23, 1896; accessed June 12, 2018.
260. *Sydney Morning Herald* 1896: Extraordinary Heat at Willcannia. <https://trove.nla.gov.au/newspaper/article/14033576?zoomLevel=5> January 18, 1896; accessed June 12, 2018.
261. National Public Radio 2010: The Heat Wave Of 1896 And The Rise Of Roosevelt. August 11, 2010 <https://www.npr.org/templates/story/story.php?storyId=129127924&ft=1&f=1022>; accessed June 12, 2018.
262. Westcott, N. 2011: The Prolonged 1954 Midwestern U.S. Heat Wave: Impacts and Responses. *American Meteorological Society* <https://doi.org/10.1175/WCAS-D-10-05002.1>; <https://journals.ametsoc.org/doi/full/10.1175/WCAS-D-10-05002>. 1 July 1, 2011; accessed June 12, 2018.
263. Becker, R. J. and Wood, R. A. 1980: 'Heat Wave'; in *Weatherwise*, 39 (4) 32-36.
264. Wikipedia: Steele, North Dakota https://en.wikipedia.org/wiki/Steele,_North_Dakota May 14, 2018; accessed June 12, 2018.
265. Suri, D. 2001: Warm Spells in September. <http://www.dandantheweatherman.com/Bereklauw/Septwarm.htm> February 16, 2001; accessed June 12, 2018.
266. Hung, T. *et al.* 2006: Assessment with satellite data of the urban heat island effects in Asian mega cities. *Internat. J. Applied Earth Observation and Geoinformation*. 8 (11): 34-48.
267. Torok, S. *et al.* 2001: Urban heat island features of southeast Australian towns. *Aust. Met. Mag.* 50 (2001): 1-13, <http://reg.bom.gov.au/amoj/docs/2001/torok.pdf> ; accessed 22 June, 2018
268. Long, E. 2010: Contiguous U.S. temperature trends using NCDC raw and adjusted data for one-per-state rural and urban station sets. *SPPI*: http://scienceandpublicpolicy.org/images/stories/papers/originals/Rate_of_Temp_Change_Raw_and_Adjusted_NCDC_Data.pdf February 27, 2010; accessed June 12, 2018.
269. Yale University Seto Lab 2018: Forecasting Urban Growth. <https://urban.yale.edu/research/theme-3> 2018; accessed June 12, 2018.
270. Watts, W. and Barrins, J. 2014: Smart urban design could save lives in future heatwaves. *The Conversation* <https://theconversation.com/smart-urban-design-could-save-lives-in-future-heatwaves-33246> October 22, 2014; accessed June 12, 2018.
271. Bedno, S.A. *et al.* 2010: Exertional Heat Illness Among Overweight U.S. Army Recruits In Basic Training. *Aviation, Space, and Environmental Medicine*, 81 (20: 107-111(5) DOI: <https://doi.org/10.3357/ASEM.2623.2010>.
272. Lisman, P. *et al.* 2014: Heat Tolerance Testing: Association Between Heat Intolerance and Anthropometric and Fitness Measurements. *AMSUS*, 179 (11): 1339-1346 <https://doi.org/10.7205/MILMED-D-14-00169>
273. Diaz, J., Garcia, R., Lopez, C., Linares, C., Tobias, A., and Prieto, L. 2005: Mortality impact of extreme winter temperatures. *International Journal of Biometeorology* 49:179–183.
274. Rooney, C., McMichael, A.J., Kovats, R.S., and Coleman, M.P. 1998: Excess mortality in England and Wales, and in Greater London, during the 1995 heat wave. *Journal of Epidemiology and Community Health* 52: 482–486.

- 
275. Laschewski, G. and Jendritzky, G. 2002: Effects of the thermal environment on human health: an investigation of 30 years of daily mortality data from SW Germany. *Climate Research* 21: 91–103.
 276. Deschenes, O. and Moretti, E. 2009: Extreme weather events, mortality, and migration. *The Review of Economics and Statistics* 91: 659–681.
 277. Robine, J-M., *et al.* 2008: “Solongo”. *Comptes Rendus Biologies*. 331 (2): 171–178. DOI:10.1016/j.crv.2007.12.001. ISSN 1631-0691. PMID 18241810.
 278. Stedman, J.R. 2004: The predicted number of air pollution related deaths in the UK during the August 2003 heatwave. *Atmospheric Environment* 38: 1087–1090.
 279. Fischer, P.H., Brunekreef, B., and Lebret, E. 2004: Air pollution related deaths during the 2003 heat wave in the Netherlands. *Atmospheric Environment* 38: 1083–1085.
 280. Fouillet, A. *et al.* 2008: Has the impact of heat waves on mortality changed in France since the European heat wave of summer 2003? A study of the 2006 heat wave. *International J Epidemiology* 37:309–317.
 281. Harlan, S. L. and Ruddell, D. M. 2011: Climate change and health in cities: impacts of heat and air pollution and potential co-benefits from mitigation and adaptation. *Current Opinion in Environmental Sustainability*, 3 (3): 126-134 <https://doi.org/10.1016/j.cosust.2011.01.001>
 282. Posey, C., 1981: 1980: Heat wave. *Weatherwise*, 33: 112–116.
 283. Goklany, I. 2007: Deaths and death rates due to extreme weather events. *Civil Society Report on Climate Change*. International Policy Network.
 284. Cary, G.J., Bradstock, R.A., Gill, A.M., Williams, R.J. 2012: *Global change and fire regimes in Australia pp149-170 in Flammable Australia: Fire regimes, biodiversity and ecosystems in a changing world*. (Eds Bradstock RA, Gill AM, Williams RJ). CSIRO Publishing, Collingwood, VIC.
 285. Russell-Smith J, Yates CP, Whitehead PJ, Smith R, Craig R 2007: Bushfires ‘Down Under’: patterns and implications of contemporary Australian landscape burning. *International Journal of Wildland Fire* 16: 361–377.
 286. Cameron, P. A. *et al.* 2009: Black Saturday: the immediate impact of the February 2009 bushfires in Victoria, Australia. *Medical Journal of Australia* 191: 11-16.
 287. Haynes, K., Handmer, J., McAneney, J. *et al.* 2010: Australian bushfire fatalities 1900–2008: exploring trends in relation to the ‘Prepare, stay and defend or leave early’ policy. *Environmental Science and Policy* 13:185-194.
 288. O’Neill S. J, and Handmer J. 2012: Responding to bushfire risk: the need for transformative adaptation. *Environmental Research Letters* 7: 14-18.
 289. Bernstein, A. S. and Rice, M. B. 2013: Lungs in a warming world: climate change and respiratory health. *CHEST Journal* 143:1455-1459.
 290. Johnston, F. 2017: Bushfires and Planned Burns – Tips for your patients in managing smoke. *Respiratory Medicine Today* 2 (2): 34-36.
 291. Johnston, F.H. *et al.* 2012: Estimated global mortality attributable to smoke from landscape fires. *Environmental Health Perspectives* 120: 695-701.
 292. Dennekamp, M., Erbas B., Sim, M., Glass, *et al.* 2011: Air pollution from bushfires and out of hospital cardiac arrests in Melbourne Australia. *Epidemiology* 22: S53.
 293. Johnston, F.H. *et al.* 2011: Extreme air pollution events from bushfires and dust storms and their association with mortality in Sydney, Australia 1994-2007. *Environmental Research* 111: 811-16.
 294. Morgan, G., Sheppard, V., Khalaj, B., Ayyar, A., Lincoln, D., Jalaludin, B. *et al.* 2010: The effects of bushfire smoke on daily mortality and hospital admissions in Sydney, Australia, 1994 to 2002. *Epidemiology* 21: 47-55.
 295. Martin, K.L. *et al.* 2013: Air pollution from bushfires and their association with hospital admissions in Sydney, Newcastle and Wollongong, Australia 1994–2007. *Australian and New Zealand Journal of Public Health* 37: 238-243.




296. Smith, H.G., Sheridan, G.J., Lane, P.N.J., Nyman, P., Haydon, S. 2011: Wildfire effects on water quality in forest catchments: A review with implications for water supply. *Journal of Hydrology* 396: 170-192.
297. Kuczera, G.A. 1985: Prediction of water yield reduction following a bushfire in Ash-mixed species eucalypt forest. *Water Supply Catchment Hydrology Research Report MMBW-W-0014*, Melbourne and Metropolitan Board of Works.
298. McFarlane A. C. 1988: The longitudinal course of posttraumatic morbidity: the range of outcomes and their predictors. *Journal of Nervous and Mental Diseases* 176: 30–9.
299. McFarlane, A.C. and Raphael, B. 1984: Ash Wednesday: the effect of a fire. *Australian and New Zealand Journal of Psychiatry* 18: 341–51.
300. Giglio, L. *et al.* 2013: Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4), *Journal of Geophysical Research-Biogeosciences* 118 (1): 317–328.
301. Hemp, A. 2005: Climate change-driven forest fires marginalise the impact of ice cap wasting on Kilimanjaro. *Global Change Biology* 11: 1013-1023.
302. Kraaij, T., Baard, J.A., Cowling, R.M., van Wilgen, B.W. 2013: Historical fire regimes in a poorly understood, fire-prone ecosystem: eastern coastal fynbos. *International Journal of Wildland Fire* 3: 277-287.
303. Bradstock, R.A., Penman, T., Boer, M., Price, O., Clarke, H. 2013: Divergent responses of fire to recent warming and drying across south-eastern Australia. *Global Change Biology* doi:10.1111/gcb.12449
304. Kasischke, E.S., Verbyla, D.L., Rupp, T.S., McGuire, A.D., Murphy, K.A., Jandt, R., *et al.* 2010: Alaska's changing fire regime: implications for the vulnerability of boreal forests. *Canadian Journal of Forest Research* 40: 1360-1370.
305. Gillett, N.P., Weaver, A.J., Zwiers, F.W., Flannigan, M.D. 2004: Detecting the effect of climate change on Canadian forest fires. *Geophysical Research Letters* 31: L18211.
306. Westerling, A.L., Hidalgo, H.G., Cayan, D.R., Swetnam, T.W. 2006: Warming and earlier spring increase Western U.S. forest wildfire activity. *Science* 313: 940-943.
307. Williams, J., *et al.* 2011: Findings and implications from a coarse-scale global assessment of recent selected megafires. Paper presented to 5th International Wildland Fire Conference, Sun City, South Africa. <http://foris.fao.org/static/pdf/fm/5thIWFFConference2011.pdf> May 9-13, 2011; accessed June 12, 2018.
308. Pausas, J. G. and Fernandez-Munoz, S. 2012: Fire regime changes in the Western Mediterranean Basin: from fuel-limited to drought-driven fire regime. *Climatic Change* 110: 215-216.
309. Chen, K. and McAneney, J. 2010: Bushfire Penetration into Urban Areas in Australia: A Spatial Analysis. Report for the Bushfire CRC. <http://www.bushfirecrc.com/sites/default/files/managed/resource/bushfire-penetration-urban-areas.pdf> January 2010; accessed June 12, 2018.
310. Hovenden, M.J. and Williams, A.L. 2010: The impacts of rising CO₂ concentrations on Australian terrestrial species and ecosystems. *Austral Ecology* 35: 665-684.
311. Flannigan, M., Cantin, A. S., de Groot, W.J., Wotton, M., Newbery, A., Gowman, L.M. 2013: Global wildland fire season severity in the 21st century. *Forest Ecology and Management* 294: 54-61.
312. Villarini, G. and Smith, J.A. 2013: Spatial and temporal variability of cloud-to-ground lightning over the continental U.S. during the period 1995–2010. *Atmospheric Research* 124: 137-148.
313. Piao, S. L. *et al.* 2013: Evaluation of terrestrial carbon cycle models for their response to climate variability and to CO₂ trends. *Glob. Change Biol.* 19: 2117–2132.
314. Price C. and Rind D. 1994: Possible implications of global climate change on global lightning distributions and frequencies. *Journal of Geophysical Research* 99 (D5): 10,823–10,831.
315. Balch, J. K, *et al.* 2017: Human-started wildfires expand the fire niche across the United States. *PNAS* 114,11, 2946-2951 <http://www.pnas.org/content/pnas/114/11/2946.full.pdf> January 6, 2017; accessed June 12, 2018.

- 
316. Steele, J. 2017: Deconstructing the Climate Demagoguery of the Wine Country Wildfire Tragedies. *Landscapes and Cycles*, <http://landscapesandcycles.net/wine-country-fires-and-climate-demogoguery.html> October 25, 2017; accessed June 12, 2018.
 317. 2009 Victorian Bushfires Royal Commission 2010: Final Report. <http://royalcommission.vic.gov.au/Commission-Reports/Final-Report/Summary/Interactive-Version.html> July 2010; accessed June 12, 2018.
 318. Queensland Government 2016: Gamba Grass. https://www.daf.qld.gov.au/__data/assets/pdf_file/0011/67466/IPA-Gamba-Grass-PP147.pdf 2016; accessed June 12, 2018.
 319. The Conversation 2017: How invasive weeds can make wildfires hotter and more frequent. https://theconversation.com/how-invasive-weeds-can-make-wildfires-hotter-and-more-frequent-89281?utm_medium=email&utm_campaign=Latest%20from%20The%20Conversation%20for%20December%2021%202017%20-%2090897694&utm_content=Latest%20from%20The%20Conversation%20f; accessed June 12, 2018.
 320. Wik, M., Pingali, P. and Broca, S. 2008: Global Agricultural Performance: Past Trends and Future Prospects, *World Bank* http://siteresources.worldbank.org/INTWDR2008/Resources/2795087-1191427986785/Pingali-Global_Agricultural_Performance.pdf; accessed June 12, 2018.
 321. Horton, G. and McMichael, T. 2008: Climate change health check 2020. *Climate Institute of Australia*. Citing: World Health Organisation. Protecting health from climate change – a toolkit for event organisers. http://www.who.int/world-health-day/toolkit/toolkit_en.pdf; accessed June 12, 2018.
 322. Heiberg, T. 2017: SA's 2017 maize crop largest on record – committee. *Moneyweb* <https://www.moneyweb.co.za/news-fast-news/sas-2017-maize-crop-largest-on-record-committee/>; accessed June 12, 2018.
 323. Intergovernmental Panel on Climate Change (IPCC) 2014: Assessing and Managing the Risks of Climate Change. https://ipcc.ch/report/ar5/wg2/docs/WGIIAR5_SPM_Top_Level_Findings.pdf accessed June 12, 2018.
 324. Asseng, S. et al. 2015: Rising temperatures reduce global wheat production. *Nature Climate Change* 5: 143–147. DOI:10.1038/nclimate2470. <https://www.nature.com/articles/nclimate2470>; accessed June 12, 2018.
 325. Stallard, B. 2014: Rising Temperatures: Wheat Production Takes a Hit in India. *Nature World News* <https://www.natureworldnews.com/articles/8208/20140723/rising-temperatures-wheat-production-takes-hit-india.htm?exe=reporter> July 23, 2014; accessed June 12, 2018.
 326. Mohan, V. 2017: Govt revises foodgrain output to record 275.68 million tonnes. *The Times of India* <https://timesofindia.indiatimes.com/india/govt-revises-foodgrain-output-to-record-275-68-million-tonnes/articleshow/60090001.cms> August 17, 2017; accessed June 12, 2018.
 327. Hillsdon, M. 2018: Wheat in heat: the 'crazy idea' that could combat food insecurity. *The Guardian* https://www.theguardian.com/global-development/2018/mar/23/heat-tolerant-durum-wheat-crazy-idea-food-insecurity?utm_source=CCNet+Newsletter&utm_campaign=e4674d7290-EMAIL_CAMPAIGN_2018_04_10&utm_medium=email&utm_term=0_fe4b2f45ef-e4674d7290-20143845 March 23, 2018; accessed June 12, 2018.
 328. Lyddon, C., 2018: Another record-breaking harvest. *World-Grain.com* http://www.world-grain.com/articles/news_home/Features/2016/11/Another_record-breaking_harves.aspx?ID=%7BF66FAB2B-AE1E-40B6-95F7-A7F92CD9B379%7D&cck=1 November 11, 2016; accessed June 12, 2018.
 329. Frich, P., Alexander, L., V., Della-Marta, P., Gleason, B., Haylock, M. et al. 2002: Observed coherent changes in climatic extremes during the second half of the twentieth century. *Climate Research* 19: 193–212.
 330. Curtin, T. 2009: Climate change and food production. *Energy and Environment* 20 (7): 1099–1116.
 331. Jablonski, L.M. et al. 2002: Plant reproduction under elevated CO₂ conditions: a meta-analysis of reports of 79 crop and wild species. *New Phytologist* 156: 9–26.
 332. Kimball, B.A., Kobayashi, K, and Bindi, M. 2002: Responses of agricultural crops to free-air CO₂ enrichment. *Advances in Agronomy* 77: 293–368.




333. Ainsworth, E. A. and Long, S. P. 2005: What have we learned from fifteen years of free-air CO₂ enrichment FACE? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytologist* 165:351-372.
334. CO₂ Coalition and New Criterion, July 2016: The Climate Surprise. Why CO₂ is good for the Earth. <http://co2coalition.org/wp-content/uploads/2016/09/The-Climate-Surprise-CO2C.pdf> accessed June 12, 2018.
335. Levine, L.H., Kasahara, H., Kopka, J., Erban, A., Fehrl, I., Kaplan, F. *et al.* 2008: Physiologic and metabolic responses of wheat seedlings to elevated and super-elevated carbon dioxide. *Advances in Space Research* 42: 1917–1928.
336. Wang, S.Y., Bunce, J.A., and Maas, J.L. 2003: Elevated carbon dioxide increases contents of antioxidant compounds in field-grown strawberries. *Journal of Agricultural and Food Chemistry* 51: 4315–4320.
337. Kimball, B.A. and Mitchell, S.T. 1981: Effects of CO₂ enrichment, ventilation, and nutrient concentration on the flavor and vitamin C content of tomato fruit. *Hort. Science* 16: 665–666.
338. Madsen, E. 1975: Effect of CO₂ environment on growth, development, fruit production and fruit quality of tomato from a physiological viewpoint. In: Chouard, P. and de Bilderling, N. (Eds.) *Phytotronics in Agricultural and Horticultural Research*. Bordas, Paris, p. 318–330.
339. Idso, S.B., Kimball, B.A., Shaw, P.E., Widmer, W., Vanderslice, J.T. *et al.* 2002: The effect of elevated atmospheric CO₂ on the vitamin C concentration of (sour) orange juice. *Agriculture, Ecosystems and Environment* 90: 1–7.
340. Caldwell, C.R., Britz, S.J., and Mirecki, R.M. 2005: Effect of temperature, elevated carbon dioxide, and drought during seed development on the isoflavone content of dwarf soybean [*Glycine max* (L.) Merrill] grown in controlled environments. *Journal of Agricultural and Food Chemistry* 53: 1125–1129.
341. Schonhof, I., Klaring, H.-P., Krumbein, A., and Schreiner, M. 2007: Interaction between atmospheric CO₂ and glucosinolates in broccoli. *Journal of Chemical Ecology* 33: 105–114.
342. Mikkelsen, M.D., Petersen, B., Olsen, C., and Halkier, B.A. 2002: Biosynthesis and metabolic engineering of glucosinolates. *Amino Acids* 22: 279–295.
343. La, G.-X., Fang, P., Teng, Y.-B., *et al.* 2009: Effect of CO₂ enrichment on the glucosinolate contents under different nitrogen levels in bolting stem of Chinese kale (*Brassica alboglabra* L.). *J. of Zhejiang University Science B* 10: 454–464.
344. Jin, C.W., Du, S.T., Zhang, Y.S., Tang, C., and Lin, X.Y. 2009: Atmospheric nitric oxide stimulates plant growth and improves the quality of spinach (*Spinacia oleracea*). *Annals of Applied Biology* 155: 113–120.
345. Gwynn-Jones, D., Jones, A.G., Waterhouse, A. *et al.* 2012: Enhanced UV-B and elevated CO₂ impacts sub-Arctic shrub berry abundance, quality and seed germination. *Ambio* 41 (Supplement 3): 256–268.
346. Goklany, I.M. 2011: Could biofuel policies increase death and disease in developing countries? *Journal of American Physicians and Surgeons* 16 (1): 9–13.
347. Carter, C. A. and Miller, H. L. 2012: Corn for Food, Not Fuel. *New York Times*, <https://www.nytimes.com/2012/07/31/opinion/corn-for-food-not-fuel.html> July 30, 2012; accessed June 12, 2018.
348. DeCicco, J.M., Liu, D.Y., Heo, J. *et al.* 2016: Carbon balance effects of U.S. biofuel production and use. *Climatic Change* 1764. DOI:10.1007/s10584-016-1764-4.
349. Fleming, A., Hobday, A. J., Farmery, A., van Putten, E. I., *et al.* 2014: Climate change risks and adaptation options across Australian seafood supply chains – A preliminary assessment. *Climate Risk Management* Vol. 1, 2014 <https://www.sciencedirect.com/science/article/pii/S2212096313000065#b0020> accessed June 12, 2018.
350. Yashodhara, B.M., Umakanth, S. *et al.* 2009: Omega-3 fatty acids: a comprehensive review of their role in health and disease. *BMJ: Postgraduate Medical Journal* 85:84-90 <http://dx.doi.org/10.1136/pgmj.2008.073338>, <http://pmj.bmj.com/content/85/1000/84> February 2009; accessed June 12, 2018.
351. Yashodhara, B.M., Umakanth, S. *et al.* 2009: Omega-3 fatty acids: a comprehensive review of their role in health and disease. *BMJ: Postgraduate Medical Journal* 85: 84-90 <http://dx.doi.org/10.1136/pgmj.2008.073338>, <http://pmj.bmj.com/content/85/1000/84> February 2009; accessed June 12, 2018.

- 
352. Ries, J.B., Cohen, A.L. and McCorkle, D.C. 2009: Marine calcifiers exhibit mixed responses to CO₂-induced ocean acidification. *Geology* 37: 1131-1134.
 353. Cheung, W.W.L, Lam, V.W.Y. *et al.* 2009: Large-scale redistribution of maximum fisheries catch potential in the global ocean under climate change. *Global Change Biology*, 16 (1): 24-25. DOI:10.1111/j.1365-2486.2009.01995.x <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1365-2486.2009.01995.x> December 2; accessed June 12, 2018.
 354. Samaila, U.R., Cheung, W.W.L, Lam, V.W.Y., Pauly, D. and Herrick, D. 2011: Climate change impacts on the biophysics and economics of world fisheries. *Nature Climate Change* 1: 449-456 doi:10.1038/nclimate1301
 355. CO₂ Coalition and *New Criterion* July 2016: The Climate Surprise. Why CO₂ is good for the Earth. <http://co2coalition.org/wp-content/uploads/2016/09/The-Climate-Surprise-CO2C.pdf> accessed June 12, 2018.
 356. Schwartz, P. and Randall, D. 2003: An Abrupt Climate Change Scenario and Its Implications for United States National Security, October 2003. <http://eesc.columbia.edu/courses/v1003/readings/Pentagon.pdf> October 2003; accessed June 12, 2018.
 357. Owain, E.L. and Maslin, M.A. 2018: Assessing the relative contribution of economic, political and environmental factors on past conflict and the displacement of people in East Africa. *Palgrave Communications*, 4:47. DOI:10.1057/s41599-018-0096-6
 358. Allen, D.W. 2011: *The Weather Makers Re-Examined*. Irenic Publications <http://irenicpublications.com.au/book-lovers/the-weather-makers-re-examined/> accessed June 12, 2018.
 359. Wikipedia: An Inconvenient Truth. https://en.wikipedia.org/wiki/An_Inconvenient_Truth June 11, 2018; accessed June 12, 2018.
 360. Tucci, J., Mitchell, J. and Goddard, D. 2007: *Children's fears, hopes and heroes: Modern Childhood in Australia*. Australian Childhood Foundation, Ringwood, Vic, National Research Centre for the Prevention of Child Abuse, Monash University, June 11, 2007, p. 5.
 361. Kiernan R. 2008: *Medical Observer*. June 13, 2008, p. 10.
 362. Oreskes, N. and Conway, E.M. 2014: Merchants of Doubt. IMbD <https://www.imdb.com/title/tt3675568/> accessed June 12, 2018.
 363. Cook, J. *et al.* 2013: Quantifying the consensus on anthropogenic global warming in the scientific literature. *Environmental Research Letters* <http://iopscience.iop.org/article/10.1088/1748-9326/8/2/024024> May 15, 2013; accessed June 12, 2018.
 364. Verheggen, B., Strengers, B., Cook, J. *et al.* 2014: Scientists' Views about Attribution of Global Warming. *Environmental Science & Technology* doi:10.1021/es501998e. <http://pubs.acs.org/doi/abs/10.1021/es501998e> July 22, 2014; accessed June 12, 2018.
 365. Strengers B, Verheggen, B and Vringer, K. 2015: Climate Science Survey. Questions and Responses, PBL Netherlands Environmental Assessment Agency http://www.pbl.nl/sites/default/files/cms/publicaties/pbl-2015-climate-science-survey-questions-and-responses_01731.pdf April 10, 2015; accessed June 12, 2018.
 366. Younger, E. 2017: Commonwealth Bank faces legal action over failure to disclose climate change risk in report. ABC. net.au <http://www.abc.net.au/news/2017-08-08/commonwealth-bank-legal-action-over-climate-change-disclosure/8786046> August 8, 2017; accessed June 12, 2018.
 367. Klimes, M. 2017: Could advisers face legal action over climate change risks? Professional Pensions <https://www.professionalspensions.com/professional-pensions/analysis/3022321/could-advisers-face-legal-action-over-climate-change-risks> December 4, 2017; accessed June 12, 2018.
 368. Johnston, I. 2017: Government facing legal action over failure to fight climate change. *Independent* <https://www.independent.co.uk/environment/climate-change-legal-action-emissions-reduction-plan-theresa-may-clientearth-a7550676.html> January 28, 2017; accessed June 12, 2018.
 369. Harrabin, R., 2017: Climate change: Ministers should be 'sued' over targets. BBC <https://www.bbc.com/news/uk-41401656> September 27, 2017; accessed June 12, 2018.




370. Ensia 2015: Are countries legally required to protect their citizens from climate change? <https://ensia.com/features/are-countries-legally-required-to-protect-their-citizens-from-climate-change/> July 15, 2015; accessed June 12, 2018.
371. Chang, C. 2016: Young people are suing governments over climate change. *News.com.au* <https://www.news.com.au/technology/environment/climate-change/young-people-are-suing-governments-over-climate-change/news-story/e327a797ab048ba2013f7f96c2d3ffbc> March 10, 2016; accessed June 12, 2018.
372. Alexander, K. 2018: Oil giant ExxonMobil counters climate-change suits by SF, other governments. *SF Gate* <https://www.sfgate.com/bayarea/article/Oil-giant-Exxon-Mobil-counters-climate-change-12483217.php> January 8, 2018; accessed June 12, 2018.
373. Leavenworth, S. 2018: Exxon Mobil takes aim at local governments. *The Lewiston Tribune* https://lmtribune.com/editors_pick/exxon-mobil-takes-aim-at-local-governments/article_e5335f12-ca7a-50d1-88bc-b1d8995f92ca.html March 5, 2018; accessed June 12, 2018.
374. Westmore, P. 2010: Tuvalu sinking? Much ado about nothing. <http://www.newsweekly.com.au/article.php?id=4236> June 26, 2010; accessed June 12, 2018.
375. Cabanes, C., Cazenave, A. and Provost, C.L. 2001: Sea level rise during the past 40 years determined from satellite and in situ observations. *Science* 294: 840-842.
376. CO₂ Coalition 2016: A *Primer on Carbon Dioxide and Climate*. <http://co2coalition.org/publications/a-primer-on-carbon-dioxide-and-climate/>; accessed June 12, 2018.
377. World Health Organization 2018: Household air pollution and health. <http://www.who.int/en/news-room/fact-sheets/detail/household-air-pollution-and-health> May 8, 2018; accessed June 12, 2018.
378. Darwall, R. 2018: The World Bank's anti-energy policy betrays its core development mission. *Washington Examiner* <https://www.washingtonexaminer.com/opinion/op-eds/the-world-banks-anti-energy-policy-betrays-its-core-development-mission> April 16, 2018; accessed June 12, 2018.
379. World Nuclear Association 2017: 'Clean Coal' Technologies, Carbon Capture & Sequestration. <http://www.world-nuclear.org/information-library/energy-and-the-environment/clean-coal-technologies.aspx> September 2017; accessed June 12, 2018.
380. World Energy Council 2018: Coal in India. <https://www.worldenergy.org/data/resources/country/india/coal/> 2018; accessed June 12, 2018.
381. Wikipedia: Mining accident. https://en.wikipedia.org/wiki/Mining_accident June 6, 2018; accessed June 12, 2018.
382. ABC News 2006: Coal Mining Steeped in History. <https://abcnews.go.com/Primetime/Mine/story?id=1475697> January 5, 2006; accessed June 12, 2018.
383. GBD 2013 Mortality and Causes of Death, Collaborators 2014: Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 385: 117–71. DOI:10.1016/S0140-6736(14)61682-2 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4340604/> December 18, 2014; accessed June 12, 2018.
384. Brook, Benedict 2017: Black lung, a potentially fatal disease that Australia eradicated decades ago, has re-emerged. *News.com.au*. <https://www.news.com.au/lifestyle/health/health-problems/black-lung-a-potentially-fatal-disease-that-australia-eradicated-decades-ago-has-reemerged/news-story/2fcc634d0261e88f4c6463d39f6a84aa> February 10, 2017; accessed June 12, 2018.
385. Joy, G. J., Colinet, J. F. and Landen, D.D. Coal workers' pneumoconiosis prevalence disparity between Australia and the United States. Centers for Disease Control and Prevention. <https://www.cdc.gov/niosh/mining/UserFiles/Works/pdfs/cwppd.pdf> January 2011; accessed June 12, 2018.
386. Suarathana, E., Laney, A.S., Storey, E., Hale, J.M., Attfield, M.D. 2011: Coal workers' pneumoconiosis in the United States: regional differences 40 years after implementation of the 1969 Federal Coal Mine Health and Safety Act. *Occup. Environ. Med.* 68(12):908-13. DOI:10.1136/oem.2010.063594.

- 
387. Queensland Government 2018: Coal dust management. <https://www.qld.gov.au/environment/pollution/monitoring/coal-dust/management>; accessed June 12, 2018.
 388. Joy, G. J., Colinet, J. F. and Landen, D.D. Coal workers' pneumoconiosis prevalence disparity between Australia and the United States. Centers for Disease Control and Prevention. <https://www.cdc.gov/niosh/mining/UserFiles/Works/pdfs/cwppd.pdf> January 2011; accessed June 12, 2018.
 389. Wikipedia: Catalytic Converter. https://en.wikipedia.org/wiki/Catalytic_converter June 9, 2018; accessed June 12, 2018.
 390. *The Times* 2017: Scientific advice promoting diesel 'was wrong'. <https://www.thetimes.co.uk/article/scientific-advice-was-wrong-6zdh9sq> April 5, 2017; accessed June 12, 2018.
 391. Jenkins, jr., Holman W. 2017: Dieselgate Is a Political Disaster. *Wall Street Journal* <https://www.wsj.com/articles/dieselgate-is-a-political-disaster-1487116586> February 14, 2017; accessed June 12, 2018.
 392. The city with the most polluted air. (Video) 2016: *The Economist* <https://www.youtube.com/watch?v=rEbNsn91XVO> January 15, 2016; accessed June 12, 2018.
 393. McGraw, Seamus 2016: Is Fracking Safe? The 10 Most Controversial Claims About Natural Gas Drilling. *Popular Mechanics*. <https://www.popularmechanics.com/science/energy/g161/top-10-myths-about-natural-gas-drilling-6386593/> May 1, 2016; accessed June 12, 2018.
 394. Harrabin, Roger 2015: Fracking: Think again, campaigner urges environmentalists <https://www.bbc.com/news/science-environment-34191713> September 10, 2015; accessed June 12, 2018.
 395. Malbran, Pia 2010: Gas Explosions Not Uncommon. *CBS Evening News* <https://www.cbsnews.com/news/gas-explosions-not-uncommon/> September 10, 2010; accessed June 12, 2018.
 396. U.S. Energy Information Administration, Frequently Asked Questions: How much carbon dioxide is produced when different fuels are burned? <https://www.eia.gov/tools/faqs/faq.php?id=73&t=11> June 8, 2018; accessed June 12, 2018.
 397. World Energy Council, World Energy Resources-Bioenergy, 2016. https://www.worldenergy.org/wp-content/uploads/2017/03/WEResources_Bioenergy_2016.pdf 2016; accessed June 12, 2018.
 398. International Energy Agency 2014: World Energy Outlook 2014 Fact sheet: Energy in sub-Saharan Africa today. https://www.iea.org/media/news/2014/press/141013_WEO_Africa_Energy_OutlookFactsheet1.pdf accessed June 12, 2018.
 399. World Health Organization 2018: Household air pollution and health. <http://www.who.int/en/news-room/fact-sheets/detail/household-air-pollution-and-health> May 8, 2018; accessed June 12, 2018.
 400. World Health Organization, 9 out of 10 people worldwide breathe polluted air. <http://www.who.int/news-room/air-pollution> ND; accessed June 12, 2018.
 401. Lelieveld, J. *et al.* 2015: The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature*, 525: 367–371.
 402. Paunio, Mikko 2018: Kicking Away the Energy Ladder, *The Global Warming Policy Foundation*, <https://www.thegwpf.org/content/uploads/2018/05/Paunio-EnergyLadder.pdf> accessed June 12, 2018.
 403. Sterman, J. D., Siegel, L. and Rooney-Varga, J.N. 2018: Does replacing coal with wood lower CO2 emissions? Dynamic life cycle analysis of wood bioenergy. *Environmental Research Letters*, 13 (1): 1-10
 404. Thurston, G.D. *et al.* 2016: Ambient particulate matter air pollution exposure and mortality in the NIH-AARP Diet and Health cohort. *Environmental Health Perspectives* 124 (4): 484-490 doi:10.1289/ehp.1509676 *Web of Science*
 405. World Health Organization, Air Pollution <http://www.who.int/airpollution/en/> accessed June 12, 2018.
 406. Tsoutsos, T. *et al.* 2005: Environmental impacts from the solar energy technologies. *Energy Policy* 33 (3):289-296 [https://doi.org/10.1016/S0301-4215\(03\)00241-6](https://doi.org/10.1016/S0301-4215(03)00241-6)
 407. Etnier, E.L. and Watson, A.P. 1981: Health and safety implications of alternative energy technologies. II. Solar. *Environmental Management* 5 (5): 409–425.



408. Bhandari, K.P. *et al.* 2015: Energy payback time (EPBT) and energy return on energy invested (EROI) of solar photovoltaic systems: A systematic review and meta-analysis. *Renewable and Sustainable Energy Reviews*. 47: 133–141.
409. PR Newswire, Solar Cells Linked to Greenhouse Gases Over 23,000 Times Worse than Carbon Dioxide According to New Book, *Green Illusions*. <http://www.digitaljournal.com/pr/738098> June 4, 2012; accessed June 12, 2018.
410. Tsai, W-T. 2008: Environmental and health risk analysis of nitrogen trifluoride (NF₃), a toxic and potent greenhouse gas. *Journal of Hazardous Materials*, 159 (2-3): 257-263 <https://doi.org/10.1016/j.jhazmat.2008.02.023>
411. Wang, B. 2008: Deaths per TWh for all energy sources: Rooftop solar power is actually more dangerous than Chernobyl. *Next Big Future* <https://www.nextbigfuture.com/2008/03/deaths-per-twh-for-all-energy-sources.html> March 14, 2008; accessed June 12, 2018.
412. Bhandari, K.P. *et al.* 2015: Energy payback time (EPBT) and energy return on energy invested (EROI) of solar photovoltaic systems: A systematic review and meta-analysis. *Renewable and Sustainable Energy Reviews*. 47: 133–141
413. Desai, J. and Nelson, M. 2017: Are we headed for a solar waste crisis? *Environmental Progress*, June 21, 2017 <http://environmentalprogress.org/big-news/2017/6/21/are-we-headed-for-a-solar-waste-crisis>; accessed June 12, 2018.
414. Desai, J. and Nelson, M. 2017: Are we headed for a solar waste crisis? *Environmental Progress*, <http://environmentalprogress.org/big-news/2017/6/21/are-we-headed-for-a-solar-waste-crisis> June 21, 2017; accessed June 12, 2018.
415. Harding, G., Harding, P., Wilkins, A. 2008: Wind turbines, flicker, and photosensitive epilepsy: Characterizing the flashing that may precipitate seizures and optimizing guidelines to prevent them. *Epilepsia* 49: 1095–1098.
416. Hoen, B., Wiser, R., Cappers, P., Thayer, M. and Sethi, G. 2011: Wind Energy Facilities and Residential Properties: The Effect of Proximity and View on Sales Prices. *Journal of Real Estate Research*. 33 (3): 279-316.
417. Gibbons, S. 2015: Gone with the wind: valuing the visual impact of wind turbines through house prices. *J. Environ Economics and Management* 72: 177-196 <https://doi.org/10.1016/j.jeem.2015.04.006>, <https://www.sciencedirect.com/science/article/pii/S0095069615000418> July 2015; accessed June 12, 2018.
418. Gibbons, S. 2015: Gone with the wind: valuing the visual impact of wind turbines through house prices. *J. Environ Economics and Management* 72: 177-196 <https://doi.org/10.1016/j.jeem.2015.04.006>, <https://www.sciencedirect.com/science/article/pii/S0095069615000418> July 2015; accessed June 12, 2018.
419. Hartman, R. S. 2017: Savoy wind turbine study is junk science. *The Berkshire Eagle*. <http://www.berkshireeagle.com/stories/letter-savoy-wind-turbine-study-is-junk-science,518890> September 8, 2017; accessed June 12, 2018.
420. Schmidt, J.H. and Klokke, M. 2014: Health Effects Related to Wind Turbine Noise Exposure: A Systematic Review. *PLOS One*. <https://doi.org/10.1371/journal.pone.0114183>, <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0114183> December 4, 2017; accessed June 12, 2018.
421. Thomsen, J., Sass, K., Odkvist, L., Arlinger, S. 2005: Local overpressure treatment reduces vestibular symptoms in patients with Meniere's disease: a clinical, randomized, multicenter, double-blind, placebo-controlled study. *Otol. Neurotol* 26: 68–73.
422. Jung, S.S., Cheung, W.S., Cheong, C., Shin, S.H. 2008: Experimental Identification of Acoustic Emission Characteristics of Large Wind Turbines with Emphasis on Infrasound and Low-Frequency Noise. *J Korean Phys Soc* 53: 1897–1905.
423. Tang, B., Wu, D., Zhao, X., Zhou, T., Zhao, W. and Wei, H. 2017: The observed impacts of wind farms on local vegetation growth in northern China. *Remote Sensing* 9: 332, doi:10.3390/rs9040332.
424. Kelly, M. J. 2017: World's First Offshore Wind Farm Retires: A Post-Mortem. *The Global Warming Policy Foundation*. https://www.thegwpf.com/worlds-first-offshore-wind-farm-retires-a-post-mortem/?utm_source=CCNet+Newsletter&utm_campaign=c4b638da74-EMAIL_CAMPAIGN_2017_10_19&utm_medium=email&utm_term=0_fe4b2f45ef-c4b638da74-20143845 October 18, 2017; accessed June 12, 2018.
425. Chang Charis 2016: Preliminary report reveals cause of South Australia blackout. *News.com.au*. <https://www.news.com.au/technology/environment/preliminary-report-reveals-cause-of-south-australia-blackout/news-story/92606772798e23e1ceec8c53f4256900> October 6, 2016; accessed June 12, 2018.

- 
426. Darwall Rupert 2018: The World Bank's anti-energy policy betrays its core development mission. *The Washington Examiner*, <https://www.washingtonexaminer.com/opinion/op-eds/the-world-banks-anti-energy-policy-betrays-its-core-development-mission> April 16, 2018; accessed June 12, 2018.
 427. Peacefmonline.com, Akosombo Dam Level Drops Low . . . Operates At Half Capacity. <http://www.peacefmonline.com/pages/local/news/201606/281562.php> June 7, 2016; accessed June 12, 2018.
 428. Shaban A. 2017: Tanzania plunged into darkness after nationwide blackout. *Africa News* <http://www.africanews.com/2017/11/30/tanzania-plunged-into-darkness-after-nationwide-blackout/> November 30, 2017; accessed June 12, 2018.
 429. MacLean L. et al. 2018: Cape Town still has water – for now. This may take political solutions as well as technical fixes. https://www.washingtonpost.com/news/monkey-cage/wp/2018/04/23/cape-town-still-has-water-for-now-this-may-take-political-solutions-as-well-as-technical-fixes/?noredirect=on&utm_term=.6552c8344813 April 23, 2018; accessed June 12, 2018.
 430. Wikipedia: Fukushima Daiichi nuclear disaster casualties. https://en.wikipedia.org/wiki/Fukushima_Daiichi_nuclear_disaster_casualties April 6, 2018; accessed June 12, 2018.
 431. Wang B. 2008: Deaths per TWh for all energy sources: Rooftop solar power is actually more dangerous than Chernobyl. *Next Big Future* <https://www.nextbigfuture.com/2008/03/deaths-per-twh-for-all-energy-sources.html> March 14, 2008; accessed June 12, 2018.
 432. Wikipedia: List of nuclear and radiation accidents by death toll. https://en.wikipedia.org/wiki/List_of_nuclear_and_radiation_accidents_by_death_toll June 14, 2018; accessed June 14, 2018.
 433. Conca, James 2012: How Deadly Is Your Kilowatt? We Rank The Killer Energy Sources. *Forbes*, <https://www.forbes.com/sites/jamesconca/2012/06/10/energys-deathprint-a-price-always-paid/#4077fbb9709b> June 10, 2012; accessed June 12, 2018.
 434. Gibbons S. 2017: Gone with the wind: Valuing the visual impacts of wind turbines through house prices. *Journal of Environmental Economics* 72, 117-196 <https://www.sciencedirect.com/science/article/pii/S0095069615000418> July 2015; accessed June 12, 2018.
 435. U. S. Nuclear Regulatory Commission, Global Laser Enrichment Facility Licensing. <https://www.nrc.gov/materials/fuel-cycle-fac/laser.html> August 2, 2017; accessed June 12, 2018.
 436. Manheimer, W. J. 2014: Fusion Breeding for Mid-Century Sustainable Power. *Journal of Fusion Energy* 33: 199. <https://doi.org/10.1007/s10894-014-9690-9>; <https://link.springer.com/article/10.1007%2Fs10894-014-9690-9> March 25, 2014; accessed June 12, 2018.
 437. Conca, James 2012: How Deadly Is Your Kilowatt? We Rank The Killer Energy Sources. *Forbes*, <https://www.forbes.com/sites/jamesconca/2012/06/10/energys-deathprint-a-price-always-paid/#4077fbb9709b> June 10, 2012; accessed June 12, 2018.
 438. IPCC 2014: *Summary for policymakers. In: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, p. 20 https://ipcc.ch/pdf/assessment-report/ar5/wg2/ar5_wgII_spm_en.pdf Accessed June 12, 2018.



CO2 Coalition Board of Directors

Jan Breslow, MD: Fredrick Henry Leonhardt Professor Rockefeller University; Head Laboratory of Biochemical Genetics and Metabolism; Senior Physician Rockefeller Hospital.

Bruce Everett, PhD: Adjunct Professor of International Business at the Georgetown School of Foreign Service; Adjunct Associate Professor of International Business at the Fletcher School, Tufts University.

Gordon Fulks, PhD: University of Chicago Laboratory for Astrophysics; Mission Research Corporation, Corbett, Oregon.

Will Happer, PhD: Cyrus Fogg Brackett Professor of Physics (emeritus) Princeton University, former Director Office of Energy Research Director of Research, U.S. Department of Energy, co-founder Magnetic Imaging Technologies.

Richard Lindzen, PhD: Emeritus Alfred P. Sloan Professor of Meteorology; member National Academy of Sciences; fellow of the American Meteorological Society and American Association for the Advancement of Science; author of 200 scientific articles and books.

Patrick Moore, PhD: Co-founder and 15-year leader of Greenpeace (1971-1986). Chairman and Chief Scientist, Ecosense Environmental. Leader, Campaign to Allow Golden Rice Now.

Rodney Nichols: former President and Chief Executive Officer of the New York Academy of Sciences; Scholar-in-Residence at the Carnegie Corporation of New York, former Executive Vice President Rockefeller University.

Norman Rogers: Founder of Rabbit Semiconductor company; policy advisor to The Heartland Institute; member of the American Geophysical Union and the American Meteorological Society.

Harrison Schmitt, PhD: geologist, astronaut and last man to walk the moon (Apollo 17), adjunct professor physics University of Wisconsin-Madison, and former U.S. Senator from New Mexico.

Leighton Steward: geologist, author, Member, The Right Climate Stuff Ex-NASA Climate Research Team.

The mission of the CO2 Coalition is to promote broader understanding of the beneficial effects of more carbon dioxide in the atmosphere around the world. The Coalition fosters informed debate on the scientific evidence, as summarized in this Primer. The Coalition's initial paper, published in the fall of 2015, urged the public to "see for yourself."



CO2 COALITION

1621 North Kent Street, Suite 603
Arlington, Virginia 22209
www.co2coalition.org
info@co2coalition.org
571-970-3180

July 2018



CO2 COALITION